



Walther  
Meißner  
Institut



BAYERISCHE  
AKADEMIE  
DER  
WISSENSCHAFTEN

Technische  
Universität  
München



# Superconductivity and Low Temperature Physics I



**Lecture Notes**  
**Winter Semester 2023/2024**

**R. Gross**  
**© Walther-Meißner-Institut**

# Chapter 4

## Microscopic Theory



Walther  
Meißner  
Institut

**BAcW**

BAYERISCHE  
AKADEMIE  
DER  
WISSENSCHAFTEN

Technische  
Universität  
München

**TUM**

# Superconductivity and Low Temperature Physics I



**Lecture No. 7**

**R. Gross**

**© Walther-Meißner-Institut**



## **4. Microscopic Theory**

### **4.1 Attractive Electron-Electron Interaction**

#### **4.1.1 Phonon Mediated Interaction**

#### **4.1.2 Cooper Pairs**

#### **4.1.3 Symmetry of Pair Wavefunction**

### **4.2 BCS Ground State**

#### **4.2.1 The BCS Gap Equation**

#### **4.2.2 Ground State Energy**

#### **4.2.3 Bogoliubov-Valatin Transformation and Quasiparticle Excitations**

### **4.3 Thermodynamic Quantities**

### **4.4 Determination of the Energy Gap**

#### **4.4.1 Specific Heat**

#### **4.4.2 Tunneling Spectroscopy**

### **4.5 Coherence Effects**

# 4. BCS Theory

- after discovery of superconductivity, initially **many phenomenological theories** have been developed
  - London theory (1935)
  - macroscopic quantum model of superconductivity (1948)
  - Ginzburg-Landau-Abrikosov-Gorkov theory (early 1950s)
- **problem:**
  - phenomenological theories do not provide insight into the microscopic processes responsible for superconductivity
  - impossible to engineer materials to increase  $T_c$ , if mechanisms are not known
- superconductivity originates from **interactions among conduction electrons**
  - theoretical models for the description of ***interacting electrons*** are required
    - very complicated: kinetic energy of conduction electrons  $\sim 5$  eV, while interaction energy  $\sim$  meV
      - ➔ find attractive interaction which causes ordering in electron system despite high kinetic energy
    - go beyond single electron (quasiparticle) models
    - not available at the time of discovery of superconductivity

# 4. BCS Theory

- development of **BCS theory** by **J. Bardeen, L.N. Cooper and J.R. Schrieffer** in 1957
  - key element is **attractive interaction** among conduction electrons
  - 1956: Cooper shows that attractive interaction results in **pair formation** and in turn in an instability of the Fermi sea of a free electron gas
  - 1957: Bardeen, Cooper and Schrieffer develop self-consistent formulation of the superconducting state: **condensation of pairs in coherent ground state**
  - paired electrons are denoted as **Cooper pairs**

- general description of interactions by exchange bosons


- Bardeen, Cooper and Schrieffer identify **phonons** as the relevant exchange bosons
- suggested by experimental observation

$$T_c \propto 1/\sqrt{M} \propto \omega_{ph} \quad \text{isotope effect}$$

- in general, detailed nature of exchange boson does not play any role in BCS theory


- many possible exchange bosons:

**magnons, polarons, plasmons, polaritons, spin fluctuations, .....**



# 4. BCS Theory

isotop effect yields hint on type of exchange boson:



data from:

E. Maxwell, Phys. Rev. 86, 235 (1952)

B. Serin, C.A. Reynolds, C. Lohman, Phys. Rev. 86, 162 (1952)

J.M. Lock, A.B. Pippard, D. Shoenberg, Proc. Cambridge Phil. Soc. 47, 811 (1951)

*in general:  $T_c \propto 1/M^{\beta^*}$*

Element	Hg	Sn	Pb	Cd	Tl	Mo	Os	Ru
Isotopen-exponent $\beta^*$	0,50	0,47	0,48	0,5	0,5	0,33	0,2	0,0

# 4.1 Attractive Electron–Electron Interaction

- **intuitive assumption:**  
superconductivity results from *ordering phenomenon of conduction electrons*
- **problem:**
  - conduction electrons have *large (Fermi) velocity* due to Pauli exclusion principle:  $\approx 10^6$  m/s  $\approx 0.01 c$
  - corresponding (Fermi) temperature is above 10 000 K
  - in contrast: transition to superconductivity occurs at  $\approx 1 - 10$  K ( $\approx$  meV)
- **task:**
  - find *interaction mechanism* that results in ordering of conduction electrons despite their high kinetic energy
  - initial attempts fail:
    - Coulomb interaction (Heisenberg, 1947)
    - magnetic interaction (Welker, 1929)
    - .....



# 4.1.1 Phonon Mediated Interaction

- known fact since 1950:
  - $T_c$  depends on isotope mass
- conclusion:
  - lattice plays an important role for superconductivity
  - initial proposals for phonon mediated e-e interaction (1950):  
*H. Fröhlich, J. Bardeen*
- **static model of lattice mediated e-e interaction:**
  - one electron causes elastic distortion of lattice:  
 attractive interaction with positive ions results in positive charge accumulation
  - second electron is attracted by this positive charge accumulation:  
 effective binding energy

intuitive picture,  
but has to be taken with care




<http://www.max-wissen.de/>

wrong suggestion:

- Cooper pairs are stable in time such as hydrogen molecule
- pairing in real space

# 4.1.1 Phonon Mediated Interaction

- *dynamic model of lattice mediated e-e interaction:*
  - moving electrons distort lattice, causing temporary positive charge accumulation along their path
    - track of positive charge cloud
    - positive charge cloud can attract second electron
  - important: positive charge cloud rapidly relaxes again → *dynamic model*



- important question: How fast relaxes positive charge cloud when electron moves through the lattice ?
- characteristic time scale  $\tau$ :
  - frequency  $\omega_q$  of lattice vibrations (phonons):  $\tau = 1/\omega_q$
  - $\omega_q \simeq 10^{12} - 10^{13}$  1/s (maximum frequency: Debye frequency  $\omega_D$ )

# 4.1.1 Phonon Mediated Interaction

- resulting range of interaction (order of magnitude estimate)
  - how far can a second electron be, to be attracted by the positive space charge before it relaxes
  - characteristic velocity of conduction electrons:  $v_F \simeq \text{few } 10^6 \text{ m/s}$
- **interaction range**:  $v_F \cdot \tau \simeq 10^6 \frac{\text{m}}{\text{s}} \cdot 10^{-13} \text{ s} \simeq 0.1 \mu\text{m}$  (is related to GL coherence length)
- important fact:
  - retarded reaction of slow ions results in large interaction range
  - **retarded interaction**
  - retarded interaction is essential for achieving attractive interaction
  - without any retardation:
    - short interaction range
    - Coulomb repulsion between electrons dominates
- retarded interaction has been addressed during discussion of screening of phonons in metals

→ **retarded interaction potential:**


$$V(\mathbf{q}, \omega) = \frac{e^2}{\epsilon(\mathbf{q}, \omega)\epsilon_0 q^2} = \underbrace{\left( \frac{e^2}{\epsilon_0(q^2 + k_s^2)} \right)}_{\text{screened Coulomb potential}} \left( 1 + \underbrace{\frac{\tilde{\Omega}_p^2(\mathbf{q})}{\omega^2 - \tilde{\Omega}_p^2(\mathbf{q})}}_{\text{correction term is negative for } \omega < \tilde{\Omega}_p(\mathbf{q}) \rightarrow \text{overscreening}} \right)$$

screened Coulomb potential  
 $1/k_s = \text{Thomas-Fermi screening length}$   
 $\sim \text{\AA}$  in metals

$$\tilde{\Omega}_p^2(\mathbf{q}) = \Omega_p^2(\mathbf{q}) / \left[ 1 + \frac{k_s^2}{q^2} \right]$$

$q$ -dependent plasma frequency of the screened ions

# 4.1.1 Phonon Mediated Interaction



$$V(\mathbf{q}, \omega) = \frac{e^2}{\epsilon(\mathbf{q}, \omega) \epsilon_0 q^2}$$

$$= \left( \frac{e^2}{\epsilon_0 (q^2 + k_s^2)} \right) \left( 1 + \frac{\tilde{\Omega}_p^2(\mathbf{q})}{\omega^2 - \tilde{\Omega}_p^2(\mathbf{q})} \right)$$

# 4.1.2 Cooper-Pairs

- **Question:** How can we formally describe the pairing interaction?
- starting point: free electron gas at  $T = 0$  (all states occupied up to  $E_F = \hbar^2 k_F^2 / 2m$ )

- **Gedanken experiment:**

- add two further electrons, which can interact via the lattice
- describe the interaction by exchange of **virtual phonon**  
virtual phonon: is generated and reabsorbed again within time  $\Delta t \lesssim 1/\omega_q$

- wave vectors of electrons after exchange of virtual phonon with wave vector  $\mathbf{q}$ :

**electron 1:**  $\mathbf{k}'_1 = \mathbf{k}_1 + \mathbf{q}$

**electron 2:**  $\mathbf{k}'_2 = \mathbf{k}_2 - \mathbf{q}$

- total momentum is conserved:  $\mathbf{K} = \mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}'_1 + \mathbf{k}'_2 = \mathbf{K}'$


- note: - since at  $T = 0$  all states are occupied below  $E_F$ , additional states have to be at  $E > E_F$

- maximum phonon energy:  $\hbar\omega_q = \hbar\omega_D$  (Debye energy)

- ➔ accessible energy interval:  $[E_F, E_F + \hbar\omega_D]$


- ➔ interaction takes place in a spherical shell with radius  $k_F$  and thickness  $\Delta k \simeq m\omega_D / \hbar k_F$

- ➔ for given  $\mathbf{K}$  only specific wave vectors  $\mathbf{k}_1, \mathbf{k}_2$  are allowed for interaction process




# 4.1.2 Cooper-Pairs

$$\mathbf{K} = \mathbf{k}_1 + \mathbf{k}_2 > 0$$



possible phase space for interaction

$$\mathbf{K} = \mathbf{k}_1 + \mathbf{k}_2 = 0$$



possible phase space is complete spherical shell

- important conclusion:** available phase space for interaction is maximum for  $\mathbf{K} = 0$  or equivalently  $\mathbf{k}_1 = -\mathbf{k}_2$

Cooper pairs with zero total momentum:  $(\mathbf{k}, -\mathbf{k})$

$$\frac{\hbar^2 k_F^2}{2m} + \hbar\omega_D = \frac{\hbar^2 (k + \Delta k)^2}{2m} \simeq \frac{\hbar^2 (k_F^2 + 2k_F \Delta k)}{2m} = \frac{\hbar^2 k_F^2}{2m} + \frac{\hbar^2 k_F \Delta k}{m} \quad \Rightarrow \quad \Delta k = \frac{m\omega_D}{\hbar k_F}$$

# 4.1.2 Cooper-Pairs

*wave function of Cooper pairs and corresponding energy eigenvalues:*

- two-particle wave function is chosen as product of two plane waves

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = a \exp(i\mathbf{k}_1 \cdot \mathbf{r}_1) \exp(i\mathbf{k}_2 \cdot \mathbf{r}_2) = a \exp(i\mathbf{k} \cdot \mathbf{r}) \quad \text{with } \mathbf{k} = \mathbf{k}_1 = -\mathbf{k}_2, \quad \mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$$

- since pair-correlated electrons are permanently scattered into new states in interval  $[k_F, k_F + \Delta k]$   
 → pair wave function = superposition of product wave functions

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{k=k_F}^{k_F+\Delta k} a_k e^{i\mathbf{k} \cdot \mathbf{r}}$$

with  $k_F < k < k_F + \Delta k$ ,  
 since restriction to energies  $E_F < E < E_F + \hbar\omega_D$

$|a_k|^2$ : probability for realization of pair  $(\mathbf{k}, -\mathbf{k})$

- note:
  - electron with  $k < k_F$  cannot participate in interaction since all states are occupied
  - we will see later that superconductor overcomes this problem by rounding-off  $f(E)$  even at  $T = 0$ 
    - superconductor first has to pay (kinetic) energy for rounding-off  $f(E)$
    - energy is obtained back by pairing interaction (potential energy)
    - net energy gain

# 4.1.2 Cooper-Pairs

*wave function of Cooper pairs and corresponding energy eigenvalues:*

- we assume that pairing interaction only depends on relative coordinate  $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$

- Schrödinger equation: 
$$-\frac{\hbar^2}{2m} (\nabla_1^2 + \nabla_2^2)\psi(\mathbf{r}_1, \mathbf{r}_2) + V(\mathbf{r})\psi(\mathbf{r}_1, \mathbf{r}_2) = E\psi(\mathbf{r}_1, \mathbf{r}_2)$$

- insert  $\psi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{k=k_F}^{k_F+\Delta k} a_k e^{i\mathbf{k}\cdot\mathbf{r}}$ , multiply by  $e^{-i\mathbf{k}'\cdot\mathbf{r}}$  and integrate over sample volume  $\Omega$

$$\int_{\Omega} \frac{\hbar^2 k^2}{m} \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV + \int_{\Omega} V(\mathbf{r}) \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV = \int_{\Omega} E \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV$$

- integration over sample volume  $\Omega$ : 
$$\int_{\Omega} \exp[i(\mathbf{k} - \mathbf{k}')\cdot\mathbf{r}] dV = \begin{cases} 0 & \text{for } \mathbf{k} \neq \mathbf{k}' \\ \Omega & \text{for } \mathbf{k} = \mathbf{k}' \end{cases}$$

$$\underbrace{\int_{\Omega} \frac{\hbar^2 k^2}{m} \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV}_{\frac{\hbar^2 k^2}{m} a_k \Omega} + \underbrace{\int_{\Omega} V(\mathbf{r}) \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV}_{\sum_{k'=k_F}^{k_F+\Delta k} a_{k'} \int V(\mathbf{r}) \exp[i(\mathbf{k} - \mathbf{k}')\cdot\mathbf{r}] dV} = \underbrace{\int_{\Omega} E \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k}\cdot\mathbf{r}) \exp[-i(\mathbf{k}'\cdot\mathbf{r})] dV}_{E a_k \Omega}$$

*scattering integral*



# 4.1.2 Cooper-Pairs

- we use abbreviation

$$V_{\mathbf{k},\mathbf{k}'} = V_{\mathbf{k}_1,\mathbf{k}_2,\mathbf{q}} = V(\mathbf{k} - \mathbf{k}') = V(\mathbf{q}) = \frac{1}{\Omega} \int V(\mathbf{r}) e^{i(\mathbf{k}-\mathbf{k}')\cdot\mathbf{r}} dV = \frac{1}{\Omega} \int V(\mathbf{r}) e^{i\mathbf{q}\cdot\mathbf{r}} dV \quad \text{with } \mathbf{k}_1 = \mathbf{k}, \mathbf{k}_2 = -\mathbf{k}, \mathbf{q} = \mathbf{k} - \mathbf{k}'$$

- result

$$\left(E - \frac{\hbar^2 k^2}{m}\right) a_k = \sum_{k'=k_F}^{k_F+\Delta k} a_{k'} V_{\mathbf{k},\mathbf{k}'}$$

**problem:**


we have to know all matrix elements  $V_{\mathbf{k},\mathbf{k}'}$  !!!

- simplifying assumption to solve the problem:

$$V_{\mathbf{k},\mathbf{k}'} = \begin{cases} -V_0 & \text{for } k' > k_F, k < k_F + \Delta k \\ 0 & \text{else} \end{cases}$$

$$\text{with } \Delta k = \frac{m\omega_D}{\hbar k_F}$$

$$\left(E - \frac{\hbar^2 k^2}{m}\right) a_k = \sum_{k'=k_F}^{k_F+\Delta k} a_{k'} V_{\mathbf{k},\mathbf{k}'} \quad \Rightarrow \quad a_k = \frac{-V_0}{E - (\hbar^2 k^2/m)} \sum_{k'=k_F}^{k_F+\Delta k} a_{k'}$$



# 4.1.2 Cooper-Pairs

- summing up over all  $k$  using  $\sum_k a_k = \sum_{k'} a_{k'}$  yields:

$$\sum_{k=k_F}^{k_F+\Delta k} a_k = V_0 \sum_{k=k_F}^{k_F+\Delta k} \frac{1}{\left(\frac{\hbar^2 k^2}{m} - E\right)} \sum_{k'=k_F}^{k_F+\Delta k} a_{k'} \quad \Rightarrow \quad 1 = V_0 \sum_{k=k_F}^{k_F+\Delta k} \frac{1}{\left(\frac{\hbar^2 k^2}{m} - E\right)}$$

- we introduce pair density of states  $\tilde{D}(E) = D(E)/2$ : sum  $\Rightarrow$  integral ( $D(E)$  = DOS for both spin directions)

$$1 = V_0 \frac{D(E_F)}{2} \int_{E_F}^{E_F + \hbar\omega_D} \frac{d\epsilon}{2\epsilon - E} \quad \text{with} \quad \epsilon = \frac{\hbar^2 k^2}{2m}$$

# 4.1.2 Cooper-Pairs

– integration and resolving for  $E$  results in

$$1 = V_0 \frac{D(E_F)}{2} \int_{E_F}^{E_F + \hbar\omega_D} \frac{d\varepsilon}{(2\varepsilon - E)} = V_0 \frac{D(E_F)}{2} \cdot \frac{1}{2} \ln|2\varepsilon - E| \Big|_{E_F}^{E_F + \hbar\omega_D} \quad \text{with } \int \frac{dx}{ax+b} = \frac{1}{a} \ln|ax + b|$$

$$\frac{4}{V_0 D(E_F)} = \ln|2E_F + 2\hbar\omega_D - E| - \ln|2E_F - E| \quad \Rightarrow \quad -\frac{4}{V_0 D(E_F)} = \ln \frac{|2E_F - E|}{|2E_F + 2\hbar\omega_D - E|}$$

$$\exp\left(-\frac{4}{V_0 D(E_F)}\right) = \frac{|2E_F - E|}{|2E_F + 2\hbar\omega_D - E|} \quad \Rightarrow \quad |2E_F + 2\hbar\omega_D - E| \exp\left(-\frac{4}{V_0 D(E_F)}\right) = |2E_F - E|$$

$$E = 2E_F - 2\hbar\omega_D \frac{\exp\left(-\frac{4}{V_0 D(E_F)}\right)}{1 - \exp\left(-\frac{4}{V_0 D(E_F)}\right)} \quad |2E_F - E| \left[1 - \exp\left(-\frac{1}{V_0 D(E_F)}\right)\right] = 2\hbar\omega_D \exp\left(-\frac{1}{V_0 D(E_F)}\right)$$

– for weak interaction  $V_0 D(E_F) \ll 1$  we obtain:

$$E \simeq 2E_F - 2\hbar\omega_D \exp\left(-\frac{4}{V_0 D(E_F)}\right)$$

- binding energy of Cooper pairs is  $\propto \hbar\omega_D$  ( $\rightarrow$  isotope effect as  $\omega_D \propto M^{-1/2}$ )
- as  $\hbar\omega_D \ll E_F$  and  $\exp\left(-\frac{4}{V_0 D(E_F)}\right) \ll 1 \rightarrow$  binding energy is very small

# 4.1.2 Cooper-Pairs

- binding energy of Cooper pairs:

$$E \simeq 2E_F - 2\hbar\omega_D \exp\left(-\frac{4}{V_0 D(E_F)}\right)$$

*important result:*

- ➔ energy of interacting electron pair is smaller than  $2E_F$
- ➔ bound pair state (Cooper pair)
- ➔ binding energy depends on  $V_0$  and maximum phonon energy  $\hbar\omega_D$

**Note 1:**

- electrons with  $k < k_F$  cannot participate in interactions as all states for  $E < E_F$  are occupied (no free scattering state)
- superconductor solves this problem by smearing out Fermi distribution even at  $T = 0$ 
  - superconductor first has to pay kinetic energy to occupy state above  $E_F$
  - increase of kinetic energy is overcompensated by pairing energy (potential energy)
  - total energy is reduced ➔ **condensation energy**

**Note 2:**

- in Gedanken experiment we have considered only two additional electrons above  $E_F$
- in real superconductor: interaction of all electrons in energy interval around  $E_F$
- electron gas becomes instable against pairing
  - ➔ instability causes transition into new ground state: **BCS ground state**

# 4.1.2 Cooper-Pairs

estimate of the interaction range from the uncertainty relation


$$\Delta k = \frac{m\omega_D}{\hbar k_F} = \frac{\omega_D}{v_F} \Rightarrow \Delta x = \frac{1}{\Delta k} = \frac{v_F}{\omega_D} \quad \text{with } v_F \sim 10^6 \text{ m/s and } \omega_D \sim 10^{13} \text{ s}^{-1} \rightarrow \text{interaction range } R \sim 100 \text{ nm}$$

how many Cooper pairs do we find in volume  $\frac{4}{3}\pi R^3$  defined by interaction range

$$\left. \begin{array}{l} \text{➤ electron density in metal: } D(E_F)/V \sim 10^{28} \text{ eV}^{-1}\text{m}^{-3} \\ \text{➤ relevant energy interval: } \hbar\omega_D \sim 0.01 - 0.1 \text{ eV} \end{array} \right\} N = 10^{28} \cdot \hbar\omega_D \cdot \frac{4}{3}\pi (10^{-7})^3 \sim 10^5 - 10^6$$

➔ strong overlap of pairs

➔ formation of *coherent many body state*



# 4.1.2 Cooper-Pairs

attractive interaction via exchange of virtual phonons: how does the matrix element  $V_{\mathbf{k},\mathbf{k}'} = V_{\mathbf{k}_1,\mathbf{k}_2,\mathbf{q}}$  look like?

- pure Coulomb interaction

$$V(\mathbf{q}) = \frac{e^2}{\epsilon_0 q^2} = \int \left( \frac{e^2}{4\pi\epsilon_0 r} \right) e^{-i\mathbf{q}\cdot\mathbf{r}} d^3r \quad \text{positive matrix element} \rightarrow \text{repulsive interaction}$$

- screened Coulomb interaction

$$V(\mathbf{q}, \omega) = \frac{e^2}{\epsilon_0(q^2 + k_s^2)} = \int \left( \frac{e^2}{4\pi\epsilon_0 r} e^{-ik_s r} \right) e^{-i\mathbf{q}\cdot\mathbf{r}} d^3r \quad \text{positive matrix element} \rightarrow \text{repulsive interaction}$$

$(k_s = \text{Thomas-Fermi wave number, } k_s \sim \pi/a)$

- screened Coulomb interaction in metals:

$$V(\mathbf{q}, \omega) = \frac{e^2}{\epsilon(\mathbf{q}, \omega)\epsilon_0 q^2} = \left( \frac{e^2}{\epsilon_0(q^2 + k_s^2)} \right) \left( 1 + \frac{\tilde{\Omega}_p^2(\mathbf{q})}{\omega^2 - \tilde{\Omega}_p^2(\mathbf{q})} \right) \quad \text{negative matrix element if } \epsilon(\mathbf{q}, \omega) < 0$$

$\rightarrow$  attractive interaction

Thomas-Fermi wave vector  $\quad q$ -dependent plasma frequency of screened ions in metal

$$\tilde{\Omega}_p^2(\mathbf{q}) = \Omega_p^2 \left[ 1 + \frac{k_s^2}{q^2} \right]$$

for small energy differences  $(E_{\mathbf{k}} - E_{\mathbf{k}'})/\hbar = \omega < \tilde{\Omega}_p(\mathbf{q})$  of the participating electrons

$\rightarrow$  **demoninator gets negative**  $\rightarrow$  attractive interaction

$\rightarrow$  cut-off frequency:  $\omega = \tilde{\Omega}_p \simeq \omega_D$  (Debye-Frequenz)

$$\Omega_p^2 = \frac{n(Ze)^2}{\epsilon_0 M}$$

# 4.1.3 Symmetry of Pair Wavefunction

What is the symmetry of the pair wavefunction?

- **important:** pair consistst of two fermions → **total wavefunction must be antisymmetric: minus sign for particle exchange**

$$\Psi(\mathbf{r}_1, \boldsymbol{\sigma}_1, \mathbf{r}_2, \boldsymbol{\sigma}_2) = \underbrace{\frac{1}{\sqrt{V}} e^{i \mathbf{K}_s \cdot \mathbf{R}_s}}_{\substack{\text{center of mass motion} \\ \text{we assume } \mathbf{K}_s = 0}} \underbrace{f(\mathbf{k}, \mathbf{r})}_{\substack{\text{orbital} \\ \text{part}}} \underbrace{\chi(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}_2)}_{\substack{\text{spin} \\ \text{part}}} = -\Psi(\mathbf{r}_2, \boldsymbol{\sigma}_2, \mathbf{r}_1, \boldsymbol{\sigma}_1)$$

$$\mathbf{R}_s = (\mathbf{r}_1 + \mathbf{r}_2)/2$$

$$\mathbf{r} = (\mathbf{r}_1 - \mathbf{r}_2)$$

$$\mathbf{K}_s = (\mathbf{k}_1 + \mathbf{k}_2)/2$$

$$\mathbf{k} = (\mathbf{k}_1 - \mathbf{k}_2)$$

- possible **spin wavefunctions**  $\chi(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}_2)$  for electron pairs

$$S = \begin{cases} 0 & m_s = 0 & \chi^a = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) & \rightarrow \text{singlet pairing, antisymmetric spin wavefunction} \\ & & & \text{symmetric orbital function: } L = 0, 2, \dots (s, d, \dots) \\ 1 & m_s = \begin{cases} -1 & \chi^s = \downarrow\downarrow \\ 0 & \chi^s = \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \\ +1 & \chi^s = \uparrow\uparrow \end{cases} & \rightarrow \text{triplet pairing, symmetric spin wavefunction} \\ & & & \text{antisymmetric orbital function: } L = 1, 3, \dots (p, f, \dots) \end{cases}$$

# 4.1.3 Symmetry of Pair Wavefunction


What is the symmetry of the pair wavefunction?

Singlet-Pairing	$S = 0$	$L = 0, 2, 4, \dots$
Triplet-Pairing	$S = 1$	$L = 1, 3, 5, \dots$


symmetric orbital wavefunction  
 antisymmetric orbital wavefunction



$L = 0$   
 s-wave  
 superconductor



$L = 1$   
 p-wave  
 superconductor



$L = 2$ :  
 d-wave  
 superconductor

- metallic superconductors:  
 $S = 0, L = 0$
- high temperature (cuprate) superconductors:  
 $S = 0, L = 2$
- suprafluid  $^3\text{He}$ :  
 $S = 1, L = 1$




# 4.1.3 Symmetry of Pair Wavefunction



- isotropic interaction:  $V_{\mathbf{k},\mathbf{k}'} = -V_0$ 
  - interaction only depends on  $|\mathbf{k}|$
  - in agreement with angular momentum  $L = 0$  (s – wave superconductor)
  - corresponding spin wavefunction must be antisymmetric
    - **spin singlet Cooper pairs ( $S = 0$ )**
  - resulting Cooper pair: 
 $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$  spin singlet Cooper pair ( $L = 0, S = 0$ )
 
    - $L = 0, S = 0$  is realized in metallic superconductors (s – wave superconductor)
    - higher orbital momentum wavefunction in cuprate superconductors (HTS):  
 $L = 2, S = 0$  (d – wave superconductor)
- **spin triplet Cooper pairs ( $S = 1$ ):**
  - realized in superfluid  $^3\text{He}$ :  $L = 1, S = 1$  (p – wave pairing)
  - evidence for  $L = 1, S = 1$  also for some heavy Fermion superconductors (e.g.  $\text{UPt}_3$ )

# 4.1.3 Symmetry of Pair Wavefunction

$L = 0, S = 0$



$L = 2, S = 0$



[Superconductivity gets an iron boost](#)


Igor I. Mazin

*Nature* **464**, 183-186(11 March 2010)

# 4.1.3 Symmetry of Pair Wavefunction








Example: iron-based superconductors – iron pnictides

pnictogens (As, P)



spacer

chalcogens (Se or Te)




- a. s-wave, e.g. in aluminium
- b. d-wave, e.g. in copper oxides
- c. two-band s-wave with the same sign, e.g. in MgB<sub>2</sub>
- d. an s<sub>±</sub>-wave, e.g. in iron-based SC

# 4.1.3 Symmetry of Pair Wavefunction


Example:  $UPt_3$

phase diagram



Michael R. Norman, *Science* 332, 196-200 (2011)


$f$ -wave ( $E_{2u}$ ) Cooper pair wavefunction in three-dimensional momentum space



# Summary of Lecture No. 7 (1)


- **microscopic theory of superconductivity**

- problem: (i) high kinetic energy of conduction electrons:  $E_{\text{kin}} \sim \text{eV}$  (corresponding to  $T \sim 10\,000\text{ K}$ )  
 (ii) small interaction strength:  $E_{\text{int}} \sim \text{meV}$  (corresponding to  $T \sim 10\text{ K}$ )  
 → find interaction resulting in ordering of conduction electrons despite high  $E_{\text{kin}}$
- Cooper (1956): even weak attractive interaction results in instability of free electron gas  
 → **pair formation: Cooper pairs**
- general description of interaction by Feynman diagram:  
 → which **exchange boson** results in attractive interaction of conduction electrons?  
 → many candidates: **phonon, magnon, polariton, plasmon, polaron, bipolaron, ...**




- **isotope effect as „smoking gun“ experiment (1951/1952)**

- transition temperature of different isotopes:  $T_c \propto 1/\sqrt{M}$   
 → as phonon frequency  $\omega_{\text{ph}} \propto 1/\sqrt{M}$  →  $T_c \propto \omega_{\text{ph}}$   
 strong evidence for attractive interaction by exchange of virtuell phonons



- **BCS-Theorie (1957)**

- qualitative discussion of attractive interaction: slow reaction of positive ions  
 → **retarded interaction**
- estimate of interaction range  $R \simeq v_F \tau \simeq v_F / \omega_D$  ( $\omega_D$  = Debye frequency)  
 $v_F \simeq 10^6 \text{ m/s}$ ,  $\omega_D \simeq 10^{13} \text{ s}^{-1}$  →  $R \simeq 100 \text{ nm}$
- $R \gg$  interaction range of screened Coulomb interaction of conduction electrons



- **attractive electron-electron interaction**

- attractive interaction via lattice vibrations (exchange of virtual phonons: Fröhlich, Bardeen)

- scattering matrix element (i) pure Coulomb interaction:  $V(\mathbf{q}) = \frac{e^2}{\epsilon_0 q^2}$  (always positive  $\rightarrow$  repulsive interaction)

- (ii) screened Coulomb interaction:  $V(\mathbf{q}, \omega) = \frac{e^2}{\epsilon(\mathbf{q}, \omega) \epsilon_0 q^2} = \left( \frac{e^2}{k_s^2 + q^2} \right) \left( 1 + \frac{\tilde{\Omega}_p^2(\mathbf{q})}{\omega^2 - \tilde{\Omega}_p^2(\mathbf{q})} \right)$

Thomas-Fermi-wave vector

$q$ -dep. plasma frequency of screened ions in metal

- for  $E_k - E_{k'} = \hbar\omega < \hbar\tilde{\Omega}_p(\mathbf{q})$  of involved electrons: denominator becomes negative

- $\rightarrow$  negative matrix element  $\rightarrow$  **attractive interaction**

- **cut-off frequency:**  $\omega = \tilde{\Omega}_p \approx \omega_D$  (Debye frequency)

- **Cooper pairs**

- „Gedanken“ experiment:

- we add 2 additional electrons to Fermi sea at  $T = 0$  and let them interact via exchange of phonons with wave number  $q$


- scattering process:
 

electron 1:	$\mathbf{k}_1 \rightarrow \mathbf{k}'_1 = \mathbf{k}_1 + \mathbf{q}$
electron 2:	$\mathbf{k}_2 \rightarrow \mathbf{k}'_2 = \mathbf{k}_2 - \mathbf{q}$
total momentum:	$\mathbf{K} = \mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}'_1 + \mathbf{k}'_2 = \mathbf{K}'$

- only states with  $E > E_F$  are accessible due to full Fermi sea

- as  $\omega_{ph} < \omega_D$ , interaction takes place in energy interval  $[E_F, E_F + \hbar\omega_D]$  corresponding to  $k_F \leq k \leq k_F + \frac{m\omega_D}{\hbar k_F} = k_F + \Delta k$

- conservation of total momentum  $\rightarrow$  wave vectors of scattering electron must be within cut surface of two intersecting circular rings of thickness  $\Delta k$   
 $\rightarrow$  **maximum cut surface (phase space) is obtained for  $\mathbf{K} = 0$  or  $\mathbf{k}_1 = -\mathbf{k}_2$**   $\rightarrow$  **Cooper pairs  $(\mathbf{k}, -\mathbf{k})$**



# Summary of Lecture No. 7 (3)

## • Cooper pair interaction

– Ansatz: pair wave function = superposition of product wave functions:  $\Psi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{k=k_F}^{k_F+\Delta k} a_k \exp(i\mathbf{k} \cdot \mathbf{r})$   $\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$

– Schrödinger equation:  $-\frac{\hbar^2}{2m}(\nabla_1^2 + \nabla_2^2) \Psi(\mathbf{r}_1, \mathbf{r}_2) + V(\mathbf{r}) \Psi(\mathbf{r}_1, \mathbf{r}_2) = E \Psi(\mathbf{r}_1, \mathbf{r}_2)$

– Vereinfachung:  $V_{\mathbf{k},\mathbf{k}'} = \begin{cases} -V_0 & \text{for } k' > k_F, k < k_F + \Delta k \\ 0 & \text{else} \end{cases}$  with  $\Delta k = \frac{m\omega_D}{\hbar k_F}$

– total energy:  $E \simeq 2E_F - 2\hbar\omega_D \exp\left(-\frac{4}{V_0 D(E_F)}\right)$  for weak interaction:  $V_0 D(E_F) \ll 1$  binding energy:  
 $E - 2E_F \propto \hbar\omega_D$  (phonon energy)

– uncertainty relation:  $\Delta k \Delta x \geq 1 \rightarrow \Delta x \leq \frac{1}{\Delta k} = \frac{v_F}{\omega_D} \simeq 100 \text{ nm}$

## • symmetry of the pair wave function

– two fermions  $\rightarrow$  **total wave function must be antisymmetric**

Singlet Pairing	$S = 0$	$L = 0, 2, 4, \dots$
Triplet Pairing	$S = 1$	$L = 1, 3, 5, \dots$

$$S = \begin{cases} 0 & m_s = 0 & \chi^a = \frac{1}{\sqrt{2}}(\uparrow\downarrow - \downarrow\uparrow) & \text{singlet pairing} \\ 1 & m_s = \begin{cases} -1 & \chi^s = \downarrow\downarrow \\ 0 & \chi^s = \frac{1}{\sqrt{2}}(\uparrow\downarrow + \downarrow\uparrow) \\ +1 & \chi^s = \uparrow\uparrow \end{cases} & & \text{triplet pairing} \end{cases}$$

– **examples:** metallic superconductors:  $S = 0, L = 0$ , high-temperature cuprate superconductors:  $S = 0, L = 2$ , superfluid  $^3\text{He}$ :  $S = 1, L = 1$



Walther  
Meißner  
Institut



BAYERISCHE  
AKADEMIE  
DER  
WISSENSCHAFTEN

Technische  
Universität  
München



# Superconductivity and Low Temperature Physics I



**Lecture No. 8**

**R. Gross**

**© Walther-Meißner-Institut**



## 4. Microscopic Theory

### 4.1 Attractive Electron-Electron Interaction

#### 4.1.1 Phonon Mediated Interaction

#### 4.1.2 Cooper Pairs

#### 4.1.3 Symmetry of Pair Wavefunction



### 4.2 BCS Ground State

#### 4.2.1 The BCS Gap Equation

#### 4.2.2 Ground State Energy

#### 4.2.3 Bogoliubov-Valatin Transformation and Quasiparticle Excitations

### 4.3 Thermodynamic Quantities

### 4.4 Determination of the Energy Gap

#### 4.4.1 Specific Heat

#### 4.4.2 Tunneling Spectroscopy

### 4.5 Coherence Effects

# 4.2 The BCS Ground State

- **discussed so far:**
  - nature of the attractive interaction
  - attractive interaction of conduction electrons by exchange of virtual phonons (only two electrons added to Fermi sea)
    - ➔ pair formation: **Cooper pair**
  - symmetry of the pair wave function
- **not yet discussed:**
  - *How does the ground state of the total electron system look like?*
  - *What is the ground state energy?*
- **we expect:**
  - pairing mechanism goes on until the Fermi sea has changed significantly
  - if pairing energy goes to zero, pairing process will stop
  - detailed theoretical description is complicated ➔ we discuss only basics

# 4.2 The BCS Ground State

*formalism of second quantization (occupation number representation) is used* (>1927, Dirac, Fock, Jordan et al.)

- 2<sup>nd</sup> quantization formalism is useful to describe quantum many-body systems
- quantum many-body states are represented in the so-called Fock (number) state basis  
→ Fock states are constructed by filling up each single-particle state with a certain number of identical particles

$$|n_1, n_2, n_3, \dots, n_\alpha, \dots\rangle \quad n_\alpha \text{ particles in state } \psi_\alpha \quad n_\alpha = \begin{cases} 0,1 & \text{fermions} \\ 0,1,2,3, \dots & \text{bosons} \end{cases}$$

- 2<sup>nd</sup> quantization formalism introduces the creation and annihilation operators to construct and handle the Fock states
- 2<sup>nd</sup> quantization formalism is also known as the canonical quantization in quantum field theory, in which the fields (wave functions of matter) are upgraded to field operators → analogous to 1<sup>st</sup> quantization, where the physical quantities are upgraded to operators

*conduction electrons can be described by wave packets*

introduction of field operators (2<sup>nd</sup> quantization of a wave function)

$$\hat{\Psi}_\sigma(\mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma} e^{i\mathbf{k}\cdot\mathbf{r}} \quad \longleftrightarrow \quad \hat{c}_\sigma(\mathbf{k}) = \hat{c}_{\mathbf{k}\sigma} = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{\Psi}_\sigma e^{-i\mathbf{k}\cdot\mathbf{r}}$$

**annihilation operator**  
(destroys state with wave number  $\mathbf{k}$ )

$$\hat{\Psi}_\sigma^\dagger(\mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}} \quad \longleftrightarrow \quad \hat{c}_\sigma^\dagger(\mathbf{k}) = \hat{c}_{\mathbf{k}\sigma}^\dagger = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{\Psi}_\sigma^\dagger e^{i\mathbf{k}\cdot\mathbf{r}}$$

**creation operator**  
(creates state with wave number  $\mathbf{k}$ )

# 4.2 The BCS Ground State

*formalism of second quantization (occupation number representation) is used* (>1927, Dirac, Fock, Jordan et al.)

**basic relations (fermionic operators):**

$$\hat{c}_{\mathbf{k}\sigma}^\dagger |0\rangle = |1\rangle \quad \hat{c}_{\mathbf{k}\sigma} |0\rangle = 0 \quad \hat{c}_{\mathbf{k}\sigma}^\dagger |1\rangle = 0 \quad \hat{c}_{\mathbf{k}\sigma} |1\rangle = |0\rangle$$

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} = n_{\mathbf{k}\sigma} \quad \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^\dagger = 1 - n_{\mathbf{k}\sigma} \quad \langle 0 | n_{\mathbf{k}\sigma} | 0 \rangle = 0; \quad \langle 1 | n_{\mathbf{k}\sigma} | 1 \rangle = 1 \quad \text{particle number operator}$$

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}^\dagger = 0 \quad \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma} = 0 \quad \text{Pauli exclusion principle}$$

**anti-commutation relations (for fermions):**

$$\{\hat{c}_{\mathbf{k}\sigma}, \hat{c}_{\mathbf{k}'\sigma'}^\dagger\} \equiv \hat{c}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}'\sigma'}^\dagger + \hat{c}_{\mathbf{k}'\sigma'}^\dagger \hat{c}_{\mathbf{k}\sigma} = \delta_{\mathbf{k}\mathbf{k}'} \delta_{\sigma\sigma'}$$

$$\{\hat{c}_{\mathbf{k}\sigma}, \hat{c}_{\mathbf{k}'\sigma'}\} = \{\hat{c}_{\mathbf{k}\sigma}^\dagger, \hat{c}_{\mathbf{k}'\sigma'}^\dagger\} = 0$$


# 4.2 The BCS Ground State

*formalism of second quantization (occupation number representation) is used* (>1927, Dirac, Fock, Jordan et al.)

pair creation and annihilation operators:

$$P_{\mathbf{k}}^{\dagger} = \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \quad \text{pair creation operator}$$

$$P_{\mathbf{k}} = \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \quad \text{pair annihilation operator}$$

$$[P_{\mathbf{k}}, P_{\mathbf{k}'}] = \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} - \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} = 0$$



the last two operators of the first term on the r.h.s. can be moved to the front by an even number of permutations → **sign is preserved**

$$[P_{\mathbf{k}}^{\dagger}, P_{\mathbf{k}'}^{\dagger}] = \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{\mathbf{k}'\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow}^{\dagger} - \hat{c}_{\mathbf{k}'\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} = 0$$

$$[P_{\mathbf{k}}, P_{\mathbf{k}'}^{\dagger}] = \delta_{\mathbf{k}\mathbf{k}'} (1 - n_{\mathbf{k}\uparrow} - n_{-\mathbf{k}\downarrow}) \quad (\text{see next slide})$$

- some of the commutator relations of the pair operators are similar to those of bosons, although the pair operators consist only of electron (fermionic) operators
- $[P_{\mathbf{k}}, P_{\mathbf{k}}^{\dagger}] \neq 0$  but not equal to  $\delta_{\mathbf{k}\mathbf{k}'}$ , as expected for bosons, depends on  $\mathbf{k}$  and  $T$
- ➔ **pair operators do commute but are not bosonic operators**

**powers of pair operators**

$$P_{\mathbf{k}}^{\dagger} P_{\mathbf{k}}^{\dagger} = (P_{\mathbf{k}}^{\dagger})^2 = \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} = \underbrace{-\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow}^{\dagger}}_{=0} \underbrace{\hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger}}_{=0} = 0$$


**antisymmetry of fermionic wavefunction requires that powers of the pair operators disappear**

# 4.2 The BCS Ground State

*formalism of second quantization (occupation number representation) is used* (>1927, Dirac, Fock, Jordan et al.)

pair creation and annihilation operators:

$$\begin{aligned}
 [P_{\mathbf{k}}, P_{\mathbf{k}}^\dagger] &= \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{-\mathbf{k}'\downarrow}^\dagger - \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{-\mathbf{k}'\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \\
 &= \hat{c}_{-\mathbf{k}\downarrow} (1 - \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow}) \hat{c}_{-\mathbf{k}'\downarrow}^\dagger - \hat{c}_{-\mathbf{k}'\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \\
 &= (1 - \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow}) \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}'\downarrow}^\dagger - \hat{c}_{-\mathbf{k}'\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \\
 &= (1 - \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow}) (1 - \hat{c}_{-\mathbf{k}'\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}) - \hat{c}_{-\mathbf{k}'\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}'\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \\
 &= \delta_{\mathbf{k}\mathbf{k}'} (1 - n_{\mathbf{k}\uparrow}) (1 - n_{-\mathbf{k}\downarrow}) - n_{-\mathbf{k}\downarrow} n_{\mathbf{k}\uparrow}
 \end{aligned}$$

$$[P_{\mathbf{k}}, P_{\mathbf{k}}^\dagger] = \delta_{\mathbf{k}\mathbf{k}'} (1 - n_{\mathbf{k}\uparrow} - n_{-\mathbf{k}\downarrow})$$

# 4.2 The BCS Ground State

formalism of second quantization (occupation number representation) is used (>1927, Dirac, Fock, Jordan et al.)

BCS Hamiltonian for  $N$  interacting electrons

$$\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{i=1}^N \left( \underbrace{-\frac{\hbar^2}{2m} \nabla_i^2}_{\text{kinetic energy}} + \underbrace{V_{\text{ext}}(\mathbf{r})}_{\text{potential energy}} \right) + \frac{1}{2} \sum_{\sigma} \sum_{i,j=1}^N \underbrace{V_{\text{int}}(\mathbf{r}_i - \mathbf{r}_j)}_{\text{interaction energy}}$$



insertion of field operators and integration over volume  $\rightarrow$  FT of  $\mathcal{H}_{\text{BCS}}$  into  $k$ -space  
(see R. Gross, A. Marx, „Festkörperphysik“, 4. Auflage, appendix H.2 or exercise sheet No. 7)

$$\mathcal{H}_{\text{BCS}} = \underbrace{\sum_{\sigma} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma}}_{\text{energy of non-interacting free electron gas}} + \underbrace{\frac{1}{2} \sum_{\sigma_1, \sigma_2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} V_{\mathbf{q}} \hat{c}_{\mathbf{k}_1 + \mathbf{q}, \sigma_1}^{\dagger} \hat{c}_{\mathbf{k}_2 - \mathbf{q}, \sigma_2}^{\dagger} \hat{c}_{\mathbf{k}_2, \sigma_2} \hat{c}_{\mathbf{k}_1, \sigma_1}}_{\text{interaction energy}}$$

energy of non-interacting free electron gas

interaction energy

operator describes scattering from state

$(\mathbf{k}_1, \sigma_1 ; \mathbf{k}_2, \sigma_2)$  into  $(\mathbf{k}_1 + \mathbf{q}, \sigma_1 ; \mathbf{k}_2 - \mathbf{q}, \sigma_2)$  by exchange of phonon with wave vector  $\mathbf{q}$

$$\varepsilon_{\mathbf{k}} = \frac{\hbar^2 \mathbf{k}^2}{2m^*}, V_{\text{ext}}(\mathbf{r}) = 0$$

$$V_{\mathbf{q}} = \frac{1}{\Omega} \int_{\Omega} V(\mathbf{r}) e^{i\mathbf{q} \cdot \mathbf{r}} dV$$

factor  $\frac{1}{2}$  avoids double counting

# 4.2 The BCS Ground State

*formalism of second quantization (occupation number representation) is used* (>1927, Dirac, Fock, Jordan et al.)

**BCS Hamiltonian for  $N$  interacting electrons**

simplification of interaction term for pairs with  $\mathbf{k}_1 = \mathbf{k}$ ,  $\mathbf{k}_2 = -\mathbf{k}$ ,  $\sigma_1 = \uparrow$ ,  $\sigma_2 = \downarrow$  and  $V_{\mathbf{q}} = V_{\mathbf{k},\mathbf{k}'}$  with  $\mathbf{q} = \mathbf{k} - \mathbf{k}'$

$$\frac{1}{2} \sum_{\sigma_1, \sigma_2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} V_{\mathbf{q}} \hat{c}_{\mathbf{k}_1 + \mathbf{q}, \sigma_1}^\dagger \hat{c}_{\mathbf{k}_2 - \mathbf{q}, \sigma_2}^\dagger \hat{c}_{\mathbf{k}_2, \sigma_2} \hat{c}_{\mathbf{k}_1, \sigma_1} \Rightarrow \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \underbrace{\hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger}_{P_{\mathbf{k}}^\dagger} \underbrace{\hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}}_{P_{\mathbf{k}'}}$$

two-particle interaction potential summation over spin yields factor 2

pair **creation** and **annihilation** operators

⇒

$$\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}$$

often the energy is given with respect to chemical potential  $\mu$

→  $\varepsilon_{\mathbf{k}} = \frac{\hbar^2 \mathbf{k}^2}{2m^*}$  is replaced by  $\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}} - \mu$



# 4.2 The BCS Ground State

*basic definitions, abbreviations, assumptions, ....*

1. weak isotropic interaction:  $V_{\mathbf{k},\mathbf{k}'} = \begin{cases} -V_0 & \text{for } |\xi_{\mathbf{k}}|, |\xi_{\mathbf{k}'}| < \hbar\omega_D \\ 0 & \text{else} \end{cases} \quad V_0 D(E_F) \ll 1$

2. pairing (Gorkov) amplitude:  $g_{\mathbf{k}\sigma_1\sigma_2} \equiv \langle c_{-\mathbf{k}\sigma_1} c_{\mathbf{k}\sigma_2} \rangle \neq 0$   
 $g_{\mathbf{k}\sigma_1\sigma_2}^* \equiv \langle c_{-\mathbf{k}\sigma_1}^\dagger c_{\mathbf{k}\sigma_2}^\dagger \rangle \neq 0$   $\langle \dots \rangle = \text{expectation value}$

3. Pauli principle: pairing amplitude is antisymmetric for interchanging spins and wave vector:

$$g_{\mathbf{k}\sigma_1\sigma_2} = -g_{-\mathbf{k}\sigma_2\sigma_1}$$

4. spin part allows to distinguish between singlet and triplet pairing:

$$S = \begin{cases} 0 & m_s = 0 & \text{singlet pairing} \\ 1 & m_s = -1, 0, +1 & \text{triplet pairing} \end{cases}$$

5. pairing potential:

$$\Delta_{\mathbf{k}\sigma_1\sigma_2} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'\sigma_1\sigma_2}$$

$$\Delta_{\mathbf{k}'\sigma_1\sigma_2}^* \equiv - \sum_{\mathbf{k}} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}\sigma_1\sigma_2}^*$$

expectation value of pairing interaction

# 4.2 The BCS Ground State

## calculation of the ground state energy

- **Hamilton operator:**

$$\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{\mathbf{k}} \underbrace{\varepsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma}}_{n_{\mathbf{k}\sigma} = \text{particle number operator}} + \sum_{\mathbf{k}, \mathbf{k}'}^N V_{\mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}$$

$$\varepsilon_{\mathbf{k}} = \frac{\hbar^2 k^2}{2m^*} = \zeta_{\mathbf{k}} + \mu$$

- how to solve the *Schrödinger equation* ?

→ most general form of  $N$ -electron wave function:

$$|\Psi_N\rangle = \sum g(\mathbf{k}_1, \dots, \mathbf{k}_l) \hat{c}_{\mathbf{k}_1\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}_1\downarrow}^{\dagger} \dots \hat{c}_{\mathbf{k}_l\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}_l\downarrow}^{\dagger} |0\rangle$$

# of possibilities to place  $N/2$  particles on  $M$  sites:

$$\frac{M!}{[M - (N/2)]! (N/2)!}$$

**problem:** huge number of possible realizations, typically  $10^{10^{20}}$

→ **mean field approach:** occupation probability of state  $\mathbf{k}$  only depends only on **average occupation probability** of other states

→ **Bardeen, Cooper** and **Schrieffer** used the following Ansatz (mean-field approach):

$$|\Psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}=\mathbf{k}_1, \dots, \mathbf{k}_M} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger}) |0\rangle$$

$|u_{\mathbf{k}}|^2$ : probability that pair state  $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$  is empty  
 $|v_{\mathbf{k}}|^2$ : probability that pair state  $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$  is occupied  
 $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$

# 4.2 The BCS Ground State

*How to guess the BCS many particle wavefunction?*

$$|\Psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}=\mathbf{k}_1, \dots, \mathbf{k}_M} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger}) |0\rangle$$

wave function assumed by Bardeen, Cooper and Schrieffer

→ assume that the macroscopic wave function  $\psi(\mathbf{r}, t) = \psi_0(\mathbf{r}, t)e^{i\theta(\mathbf{r}, t)}$  can be described by a **coherent many particle state of fermions** (motivated by strong overlap of Cooper pairs)

- **coherent state of bosons**

discussed first by **Erwin Schrödinger** in 1926 when searching for a state of the quantum mechanical harmonic oscillator approximating best the behavior of a classical harmonic oscillator

E. Schrödinger, Der stetige Übergang von der Mikro- zur Makromechanik, *Die Naturwissenschaften* 14, 664-666 (1926).

transferred later by **Roy J. Glauber** to Fock state representation

R. J. Glauber, Coherent and Incoherent States of the Radiation Field, *Phys. Rev.* 131, 2766-2788 (1963).

Nobel Prize in Physics 2005 "**for his contribution to the quantum theory of optical coherence**", with the other half shared by John L. Hall and Theodor W. Hänsch.

# 4.2 The BCS Ground State

- **Fock state representation of coherent state of bosons** (e.g. laser light)

**coherent state  $|\alpha\rangle$**  is expressed as an infinite linear combination of number (Fock) states

$$|\phi_n\rangle = \frac{1}{\sqrt{n!}} (a^\dagger)^n |0\rangle$$

boson creation operator  $\nearrow$   $\nwarrow$  vacuum state

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |\phi_n\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{(\alpha a^\dagger)^n}{n!} |0\rangle = e^{-|\alpha|^2/2} e^{(\alpha a^\dagger)} |0\rangle$$

**Schrödinger (1926)**

normalization

$\alpha = |\alpha|e^{i\varphi}$  is complex number

probability for occupation of  $n$  particles is given by Poisson distribution



$$P(n) = |\langle \phi_n | \alpha \rangle|^2 = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2} = \frac{\langle n \rangle^n}{n!} e^{-\langle n \rangle} \quad \langle n \rangle = |\alpha|^2 = \bar{N}$$

- expectation value of number operator:  $\langle n \rangle = \bar{N} = |\alpha|^2, \quad (\Delta N)^2 = \text{var}(n) = |\alpha|^2 = \bar{N} \gg 1$
- relative standard deviation:  $\frac{\Delta N}{\bar{N}} = \frac{1}{\sqrt{\bar{N}}} \ll 1 \quad (\text{as } \bar{N} \gg 1)$
- uncertainty relation  $\Delta N \Delta \varphi \geq \frac{1}{2}, \quad \Delta \varphi \ll 1$

application: **coherent photonic state generated by laser**

# 4.2 The BCS Ground State

- *Poisson distribution*




# 4.2 The BCS Ground State

- *Poisson and normal distribution*

$$P(n) = |\langle \phi_n | \alpha \rangle|^2 = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2}$$

for large  $\bar{N} = |\alpha|^2$  the Poisson distribution approaches the normal (Gaussian) distribution:

$$P_{\bar{N}}(n) = \frac{1}{\sqrt{2\pi\bar{N}}} \exp\left(-\frac{(n - \bar{N})^2}{2\bar{N}}\right)$$



# 4.2 The BCS Ground State

- Fock state representation of *coherent state of fermions*

starting point: coherent bosonic state

$$|\alpha\rangle = e^{-|\alpha|^2/2} \exp(\alpha a^\dagger) |0\rangle$$

in analogy: **coherent fermionic state**

$$|\Psi_{\text{BCS}}\rangle = c_1 \exp\left(\sum_{\mathbf{k}} \alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger\right) |0\rangle$$

summation over  $\mathbf{k}$  since we have many fermionic modes

- we make use of the fact that higher powers of fermionic creation operators disappear due to **Pauli principle** (key difference to bosonic system):

$$P_{\mathbf{k}}^\dagger P_{\mathbf{k}}^\dagger = (P_{\mathbf{k}}^\dagger)^2 = \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger = -\hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger = 0$$

$$\Rightarrow |\Psi_{\text{BCS}}\rangle = c_1 \exp\left(\sum_{\mathbf{k}} \alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger\right) |0\rangle = c_1 \prod_{\mathbf{k}} \exp(\alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger) |0\rangle = c_1 \prod_{\mathbf{k}} (1 + \alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger) |0\rangle$$

**normalization:**  $\langle \Psi_{\text{BCS}}^* | \Psi_{\text{BCS}} \rangle = c_1^2 \langle 0 | \prod_{\mathbf{k}} (1 + \alpha_{\mathbf{k}}^* P_{\mathbf{k}}) (1 + \alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger) |0\rangle = 1$  satisfied if all factors = 1

$$1 = c_1^2 \langle 0 | (1 + \alpha_{\mathbf{k}}^* P_{\mathbf{k}}) (1 + \alpha_{\mathbf{k}} P_{\mathbf{k}}^\dagger) |0\rangle = c_1^2 (1 + |\alpha_{\mathbf{k}}|^2)$$



$$c_1 = \frac{1}{\sqrt{(1 + |\alpha_{\mathbf{k}}|^2)}}$$

# 4.2 The BCS Ground State

- BCS ground state as *coherent state of fermions*

$$|\Psi_{\text{BCS}}\rangle = c_1 \prod_{\mathbf{k}} (1 + \alpha_{\mathbf{k}} P_{\mathbf{k}}^{\dagger}) |0\rangle = \frac{1}{\sqrt{(1 + |\alpha_{\mathbf{k}}|^2)}} \prod_{\mathbf{k}} (1 + \alpha_{\mathbf{k}} P_{\mathbf{k}}^{\dagger}) |0\rangle$$



$$|\Psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger}) |0\rangle$$

$$\left. \begin{aligned} u_{\mathbf{k}} &= \frac{1}{\sqrt{(1 + |\alpha_{\mathbf{k}}|^2)}} \\ v_{\mathbf{k}} &= \frac{\alpha_{\mathbf{k}}}{\sqrt{(1 + |\alpha_{\mathbf{k}}|^2)}} \end{aligned} \right\}$$

coherence factors

*coherent superposition of pair states* → *only average pair number is fixed*

$$\Delta N = \sqrt{\bar{N}} \gg 1 \quad \frac{\Delta N}{\bar{N}} = \frac{1}{\sqrt{\bar{N}}} \ll 1 \quad \Delta N \Delta \varphi \geq \frac{1}{2} \Rightarrow \Delta \varphi \ll 1$$

→ uncertainties  $\Delta N/\bar{N}$  and  $\Delta \varphi/2\pi$  are very small for large average pair number  $\bar{N}$

$u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  are complex probability amplitudes:

$|u_{\mathbf{k}}|^2$ : probability that pair state ( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is empty  
 $|v_{\mathbf{k}}|^2$ : probability that pair state ( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is occupied  
 $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$



# 4.2 The BCS Ground State

- *some expectation values (1):*

$$|\Psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}=\mathbf{k}_1, \dots, \mathbf{k}_M} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger) |0\rangle$$

we use the identities:

- $\langle 0|\phi|\Psi\rangle = \langle \phi|0\rangle \langle \Psi|0\rangle$
- $\langle \phi|(\mathcal{A}\mathcal{B})^\dagger|\Psi\rangle = \langle \phi|\mathcal{B}^\dagger\mathcal{A}^\dagger|\Psi\rangle$

single spin particle number

$$\langle n_{\mathbf{k}\uparrow} \rangle = \langle \Psi_{\text{BCS}} | \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} | \Psi_{\text{BCS}} \rangle$$

$$= \langle 0 | (u_{\mathbf{k}}^* + v_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow}) \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger) \times \underbrace{\prod_{\mathbf{l} \neq \mathbf{k}} (u_{\mathbf{l}}^* + v_{\mathbf{l}}^* \hat{c}_{-\mathbf{l}\downarrow} \hat{c}_{\mathbf{l}\uparrow}) (u_{\mathbf{l}} + v_{\mathbf{l}} \hat{c}_{\mathbf{l}\uparrow}^\dagger \hat{c}_{-\mathbf{l}\downarrow}^\dagger)}_{|u_{\mathbf{l}}|^2 + |v_{\mathbf{l}}|^2 = 1} |0\rangle$$

$$\langle n_{\mathbf{k}\uparrow} \rangle = |u_{\mathbf{k}}|^2 \underbrace{\langle 0 | \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} | 0 \rangle}_{=0} + u_{\mathbf{k}}^* v_{\mathbf{k}} \underbrace{\langle 0 | \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger | 0 \rangle}_{=0} + v_{\mathbf{k}}^* u_{\mathbf{k}} \underbrace{\langle 0 | \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} | 0 \rangle}_{=0} + |v_{\mathbf{k}}|^2 \underbrace{\langle 0 | \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger | 0 \rangle}_{=1}$$

since  $n_{\mathbf{k}\uparrow}|0\rangle = 0|0\rangle$  use even number of permutations to transform into terms  $n_{\mathbf{k}\uparrow}|0\rangle = 0|0\rangle$

use anti-commutator  $\{\hat{c}_{\mathbf{k}\sigma}, \hat{c}_{\mathbf{k}'\sigma'}^\dagger\} = \delta_{\mathbf{k}\mathbf{k}'} \delta_{\sigma\sigma'}$

$$\langle n_{\mathbf{k}\uparrow} \rangle = |v_{\mathbf{k}}|^2$$

average total pair number

$$\bar{N} = \langle \mathcal{N} \rangle = \sum_{\mathbf{k}\sigma} \langle n_{\mathbf{k}\sigma} \rangle = \sum_{\mathbf{k}\sigma} |v_{\mathbf{k}}|^2 = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2 = \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2 - |u_{\mathbf{k}}|^2 + 1$$

(see exercise sheet No. 7 for detailed derivation)

# 4.2 The BCS Ground State

- *some expectation values (2a):*

(see exercise sheet No. 7 for detailed derivation)

statistical fluctuation of average particle number

$$(\Delta N)^2 = \langle \mathcal{N} - \langle \mathcal{N} \rangle \rangle^2 = \langle \mathcal{N}^2 \rangle - \langle \mathcal{N} \rangle^2$$

$$\bar{N} = \langle \mathcal{N} \rangle = \sum_{\mathbf{k}\sigma} \langle n_{\mathbf{k}\sigma} \rangle = \sum_{\mathbf{k}\sigma} |v_{\mathbf{k}}|^2 = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2$$

$$(\Delta N)^2 = \left\langle \left( \sum_{\mathbf{k},\sigma} n_{\mathbf{k}\sigma} \right)^2 \right\rangle - \left( \left\langle \sum_{\mathbf{k},\sigma} n_{\mathbf{k}\sigma} \right\rangle \right)^2 = 2 \sum_{\mathbf{k},\mathbf{k}'} \langle n_{\mathbf{k}} n_{\mathbf{k}'} \rangle - 2 \sum_{\mathbf{k},\mathbf{k}'} \langle n_{\mathbf{k}} \rangle \langle n_{\mathbf{k}'} \rangle$$

- i.  $\mathbf{k} \neq \mathbf{k}'$  we obtain contributions  $\langle n_{\mathbf{k}\sigma} n_{\mathbf{k}'\sigma} \rangle$  and  $\langle n_{\mathbf{k}\sigma} \rangle \langle n_{\mathbf{k}'\sigma} \rangle$ ; since there is no correlation between the calculation of  $\sum_{\mathbf{k}}$  and  $\sum_{\mathbf{k}'}$  we obtain  $\langle n_{\mathbf{k}\sigma} n_{\mathbf{k}'\sigma} \rangle = \langle n_{\mathbf{k}\sigma} \rangle \langle n_{\mathbf{k}'\sigma} \rangle$

→ the contributions  $\langle n_{\mathbf{k}\sigma} n_{\mathbf{k}'\sigma} \rangle$  and  $\langle n_{\mathbf{k}\sigma} \rangle \langle n_{\mathbf{k}'\sigma} \rangle$  just cancel each other in the sums due to the minus sign

- ii.  $\mathbf{k} = \mathbf{k}'$  we use  $\langle n_{\mathbf{k}\sigma}^2 \rangle = \langle c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} \rangle = \langle c_{\mathbf{k}\sigma}^\dagger (1 - c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}) c_{\mathbf{k}\sigma} \rangle = \underbrace{\langle c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} \rangle}_{=|v_{\mathbf{k}}|^2} - \underbrace{\langle c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma} \rangle}_{=-|v_{\mathbf{k}}|^4} = |v_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^4$

$$(\Delta N)^2 = 2 \sum_{\mathbf{k}} \langle n_{\mathbf{k}} n_{\mathbf{k}} \rangle - \sum_{\mathbf{k}} \langle n_{\mathbf{k}} \rangle \langle n_{\mathbf{k}} \rangle = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^4 - 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^4 = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2$$

# 4.2 The BCS Ground State

- *some expectation values (2b):*

(see exercise sheet No. 7 for detailed derivation)

statistical fluctuation of average particle number

$$(\Delta N)^2 = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2$$

$$\Delta N = \sqrt{2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2} = \sqrt{\bar{N}}$$

$\Delta N$  gets very large for large  $\bar{N}$ , but **relative fluctuation  $\Delta N / \bar{N}$  becomes vanishingly small !!**

# 4.2 The BCS Ground State

- *some expectation values (3):*

(see exercise sheets for detailed derivation)

pairing or Gorkov amplitude

$$g_{\mathbf{k}\sigma_1\sigma_2} \equiv \langle \Psi_{\text{BCS}} | c_{-\mathbf{k}\sigma_1} c_{\mathbf{k}\sigma_2} | \Psi_{\text{BCS}} \rangle = u_{\mathbf{k}} v_{\mathbf{k}}^*$$

$$g_{\mathbf{k}\sigma_1\sigma_2}^\dagger \equiv \langle \Psi_{\text{BCS}} | c_{-\mathbf{k}\sigma_1}^\dagger c_{\mathbf{k}\sigma_2}^\dagger | \Psi_{\text{BCS}} \rangle = u_{\mathbf{k}}^* v_{\mathbf{k}}$$


BCS Hamiltonian

$$\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}$$

$$\langle \Psi_{\text{BCS}} | \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} | \Psi_{\text{BCS}} \rangle = |v_{\mathbf{k}}|^2$$

$$\langle \Psi_{\text{BCS}} | \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} | \Psi_{\text{BCS}} \rangle = v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*$$

$$\langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} | \Psi_{\text{BCS}} \rangle = \underbrace{2 \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} |v_{\mathbf{k}}|^2}_{= \bar{N} \varepsilon_{\mathbf{k}} \text{ (kinetic energy)}} + \underbrace{\sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*}_{\text{interaction energy}}$$



# 4.2 The BCS Ground State

*task: find the minimum of the expectation value  $\langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} | \Psi_{\text{BCS}} \rangle$  by variational method (@  $T = 0$ )*

we take the energy relative to the chemical potential  $\mu$

$$\langle E_{\text{BCS}} - \bar{N} \mu \rangle = \langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} - \bar{N} \mu | \Psi_{\text{BCS}} \rangle = 2 \sum_{\mathbf{k}} (\xi_{\mathbf{k}} + \mu) |v_{\mathbf{k}}|^2 - \bar{N} \mu + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*$$

$$\langle E_{\text{BCS}} - \bar{N} \mu \rangle = \langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} - \bar{N} \mu | \Psi_{\text{BCS}} \rangle = 2 \sum_{\mathbf{k}} \xi_{\mathbf{k}} |v_{\mathbf{k}}|^2 - \bar{N} \mu + \bar{N} \mu + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*$$

$$\varepsilon_{\mathbf{k}} = \frac{\hbar^2 k^2}{2m^*} = \xi_{\mathbf{k}} + \mu$$

$$\bar{N} = 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2$$

$$\delta \left\{ 2 \sum_{\mathbf{k}} \xi_{\mathbf{k}} |v_{\mathbf{k}}|^2 + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^* \right\} = 0$$

**minimization of expectation value by variation of the probability amplitudes yields expressions for  $|u_{\mathbf{k}}|^2$  and  $|v_{\mathbf{k}}|^2$**

# 4.2 The BCS Ground State

**Method 1:** we assume that  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  are real and satisfy  $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$  (Bardeen, Cooper, Schrieffer: 1957)

$$u_{\mathbf{k}} = \sin \theta_{\mathbf{k}}, \quad v_{\mathbf{k}} = \cos \theta_{\mathbf{k}}, \quad \text{and} \quad 2 \sin \theta_{\mathbf{k}} \cos \theta_{\mathbf{k}} = \sin 2\theta_{\mathbf{k}}$$

$$\langle E_{\text{BCS}} - \bar{N} \mu \rangle = 2 \sum_{\mathbf{k}} \xi_{\mathbf{k}} |v_{\mathbf{k}}|^2 + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^* = 2 \sum_{\mathbf{k}} \xi_{\mathbf{k}} \cos^2 \theta_{\mathbf{k}} + \frac{1}{4} \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \sin 2\theta_{\mathbf{k}} \sin 2\theta_{\mathbf{k}'}$$

**minimization**  $\frac{\partial \langle E_{\text{BCS}} - \bar{N} \mu \rangle}{\partial \theta_{\mathbf{l}}} = 0$

$$\frac{\partial \langle E_{\text{BCS}} - \bar{N} \mu \rangle}{\partial \theta_{\mathbf{l}}} = 0 = 2\xi_{\mathbf{l}} \underbrace{(-2 \cos \theta_{\mathbf{l}} \sin \theta_{\mathbf{l}})}_{= -\sin 2\theta_{\mathbf{l}}} + \underbrace{\frac{1}{4} \frac{\partial}{\partial \theta_{\mathbf{l}}} \sum_{\mathbf{k}} \sin 2\theta_{\mathbf{k}} \sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \sin 2\theta_{\mathbf{k}'}}_{\frac{1}{4} (2 \cos 2\theta_{\mathbf{l}}) \sum_{\mathbf{k}'} V_{\mathbf{l}, \mathbf{k}'} \sin 2\theta_{\mathbf{k}'} + \sum_{\mathbf{k}} \sin 2\theta_{\mathbf{k}} V_{\mathbf{k}, \mathbf{l}} 2 \cos 2\theta_{\mathbf{l}}}$$

$$2\xi_{\mathbf{l}} \sin 2\theta_{\mathbf{l}} = \frac{1}{2} \cos 2\theta_{\mathbf{l}} \left( \sum_{\mathbf{k}'} V_{\mathbf{l}, \mathbf{k}'} \sin 2\theta_{\mathbf{k}'} + \sum_{\mathbf{k}} \sin 2\theta_{\mathbf{k}} V_{\mathbf{k}, \mathbf{l}} \right) \stackrel{V_{\mathbf{l}, \mathbf{k}'} = V_{\mathbf{k}, \mathbf{l}}}{=} \cos 2\theta_{\mathbf{l}} \sum_{\mathbf{k}'} V_{\mathbf{l}, \mathbf{k}'} \sin 2\theta_{\mathbf{k}'}$$

$$\tan 2\theta_{\mathbf{l}} = \frac{\sum_{\mathbf{k}'} V_{\mathbf{l}, \mathbf{k}'} \sin \theta_{\mathbf{k}'} \cos \theta_{\mathbf{k}'}}{\xi_{\mathbf{l}}}$$

# 4.2 The BCS Ground State

- we switch back to old summation ( $\mathbf{l} \rightarrow \mathbf{k}$ ) and restore  $u_{\mathbf{k}} = \sin \theta_{\mathbf{k}}$ ,  $v_{\mathbf{k}} = \cos \theta_{\mathbf{k}}$ :

$$\Rightarrow \tan 2\theta_{\mathbf{k}} = \frac{\sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} u_{\mathbf{k}'} v_{\mathbf{k}'}}{\xi_{\mathbf{k}}}$$

- we further use the pairing strength  $\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} u_{\mathbf{k}'} v_{\mathbf{k}'}$   $\Rightarrow \tan 2\theta_{\mathbf{k}} = - \frac{\Delta_{\mathbf{k}}}{\xi_{\mathbf{k}}}$

- with  $\tan 2\theta_{\mathbf{k}} = \frac{\sin 2\theta_{\mathbf{k}}}{\cos 2\theta_{\mathbf{k}}} = \frac{2 \sin \theta_{\mathbf{k}} \cos \theta_{\mathbf{k}}}{\underbrace{\cos^2 \theta_{\mathbf{k}}}_{\frac{1}{2} + \frac{1}{2} \cos 2\theta_{\mathbf{k}}} - \underbrace{\sin^2 \theta_{\mathbf{k}}}_{\frac{1}{2} - \frac{1}{2} \cos 2\theta_{\mathbf{k}}}} = \frac{2u_{\mathbf{k}}v_{\mathbf{k}}}{u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2}$  we obtain  $\Rightarrow \frac{2u_{\mathbf{k}}v_{\mathbf{k}}}{u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2} = - \frac{\Delta_{\mathbf{k}}}{\xi_{\mathbf{k}}}$

- we define  $E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2}$  and obtain the following expressions for  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  minimizing the energy

$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$|v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$$

$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

for  $k$ -independent  $\Delta_{\mathbf{k}}$ : minimum energy is  $E_{\mathbf{k}} = \Delta$

we will see later that  $E_{\mathbf{k}}$  is the energy required to add a single excitation to the ground state

$\rightarrow$  **minimum excitation energy is required, therefore  $\Delta$  represents an energy gap in the excitation spectrum**

**pairing amplitude**

**self-consistent gap equation**

# 4.2 The BCS Ground State

$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$|v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$$

$|v_{\mathbf{k}}|^2$ : probability that  $\mathbf{k}$  is occupied

→ probability  $|v_{\mathbf{k}}|^2$  is smeared out around Fermi level even at  $T = 0$ :


*increase of kinetic energy*

→ smearing is required to allow for pairing interaction:

*reduction of potential energy >*

*increase of kinetic energy*

→  $|v_{\mathbf{k}}|^2 \simeq f(T = T_c)$





# 4.2 The BCS Ground State

**Method 2:** we use the method of **Lagrangian multipliers**

- we use the following two constraints:
 
$$\phi_1 = 0 = \langle \mathcal{N} \rangle - 2 \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2 = \langle \mathcal{N} \rangle - \sum_{\mathbf{k}} |v_{\mathbf{k}}|^2 - |u_{\mathbf{k}}|^2 + 1$$

$$\phi_2 = 0 = |u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 - 1 = u_{\mathbf{k}} u_{\mathbf{k}}^* + v_{\mathbf{k}} v_{\mathbf{k}}^* - 1$$

$$\mathcal{L}(u_{\mathbf{k}}^*, v_{\mathbf{k}}^*, \lambda_1, \lambda_2) = \langle E_{\text{BCS}} \rangle - \lambda_1 \phi_1 - \lambda_2 \phi_2 \quad \lambda_1, \lambda_2: \text{Lagrangian multipliers}$$

$$\text{with } \langle E_{\text{BCS}} \rangle = 2 \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} |v_{\mathbf{k}}|^2 + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}} u_{\mathbf{k}'}^* = 2 \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} (|v_{\mathbf{k}}|^2 - |u_{\mathbf{k}}|^2 + 1) + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}} u_{\mathbf{k}'}^*$$

- by setting the partial derivative of the Lagrangian function  $\mathcal{L}$  with respect to  $u_{\mathbf{k}}^*$  and  $v_{\mathbf{k}}^*$  to zero we obtain the eigenvalue eqns:

$$\begin{aligned} (\varepsilon_{\mathbf{k}} - \lambda_1) u_{\mathbf{k}} + \Delta_{\mathbf{k}} v_{\mathbf{k}} - \lambda_2 u_{\mathbf{k}} &= 0 \\ \Delta_{\mathbf{k}}^\dagger u_{\mathbf{k}} - (\varepsilon_{\mathbf{k}} - \lambda_1) v_{\mathbf{k}} - \lambda_2 v_{\mathbf{k}} &= 0 \end{aligned} \quad \Rightarrow \quad \begin{pmatrix} (\varepsilon_{\mathbf{k}} - \lambda_1) & \Delta_{\mathbf{k}} \\ \Delta_{\mathbf{k}}^\dagger & -(\varepsilon_{\mathbf{k}} - \lambda_1) \end{pmatrix} \begin{pmatrix} u_{\mathbf{k}} \\ v_{\mathbf{k}} \end{pmatrix} = \lambda_2 \begin{pmatrix} u_{\mathbf{k}} \\ v_{\mathbf{k}} \end{pmatrix}$$

$$\text{with } \Delta_{\mathbf{k}} = -\sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} g_{\mathbf{k}'} = -\sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} u_{\mathbf{k}'} v_{\mathbf{k}'}^*$$

- physical meaning of the Lagrangian multipliers
  - $\lambda_1$  shifts the energy and corresponds to the chemical potential  $\mu$
  - $\lambda_2$  corresponds to the eigenvalue of the vector  $(u_{\mathbf{k}}, v_{\mathbf{k}})$  and is given by the energy  $\pm E_{\mathbf{k}}$  of the quasiparticles excited out of the condensate

- solving the eigenvalue eqns yields  $E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2}$ ,  $|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$ ,  $|v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$ ,  $u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$

# 4.2.1 The BCS Gap Equation

solution of the self-consistent gap equation ( $T = 0$ )

$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}} = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2\sqrt{\xi_{\mathbf{k}'}^2 + |\Delta_{\mathbf{k}'}|^2}}$$

cannot be solved analytically in the general case

- simple solution only if the gap  $\Delta_{\mathbf{k}}$  and the interaction potential  $V_{\mathbf{k},\mathbf{k}'}$  are assumed  $\mathbf{k}$ -independent:  $\Delta_{\mathbf{k}} = \Delta$ ,  $V_{\mathbf{k},\mathbf{k}'} = -V_0$

$$1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2\sqrt{\xi_{\mathbf{k}'}^2 + |\Delta|^2}} \xrightarrow[\text{with pair density } \tilde{D}(E) \approx D(E_F)/2]{\text{transforming sum into integration}} 1 = \frac{V_0 D(E_F)}{2} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{d\xi}{2\sqrt{\xi^2 + |\Delta|^2}}$$

- with  $\int \frac{dx}{\sqrt{x^2+a^2}} = \text{arcsinh}\left(\frac{x}{a}\right)$  we obtain

$$1 = \frac{V_0 D(E_F)}{4} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{d\xi}{\sqrt{\xi^2 + |\Delta|^2}} = \frac{V_0 D(E_F)}{4} \text{arcsinh}\left(\frac{\hbar\omega_D}{\Delta}\right) \Big|_{-\hbar\omega_D}^{+\hbar\omega_D} = \frac{V_0 D(E_F)}{2} \text{arcsinh}\left(\frac{\hbar\omega_D}{\Delta}\right)$$

$$\Delta = \frac{\hbar\omega_D}{\sinh(2/V_0 D(E_F))} \approx 2\hbar\omega_D e^{-2/V_0 D(E_F)}$$

energy gap corresponds to binding energy estimated for single Cooper pair

factor 2 in argument of exp. function is missing, since we have assumed that the two additional electrons are in the interval  $[E_F - \hbar\omega_D, E_F + \hbar\omega_D]$  and not in  $[E_F, E_F + \hbar\omega_D]$  as assumed previously in „Gedanken“ experiment

$V_0 D(E_F) \ll 1$ : weak coupling approximation,  $\sinh x \approx \frac{1}{2} \exp x$

# 4.2.2 Ground State Energy

## calculation of the BCS condensation energy

(see exercise sheet No. 8 for detailed derivation)

- calculate expectation value of BCS Hamiltonian for  $T = 0$

$$E_{\text{BCS}} = \langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} - \mu \mathcal{N} | \Psi_{\text{BCS}} \rangle = 2 \sum_{\mathbf{k}} \xi_{\mathbf{k}} |v_{\mathbf{k}}|^2 + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*$$

energy relative to chemical potential

$$\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}} - \mu$$

- we plug in the results for the coherence factors and the pair amplitude

$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right], \quad |v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right], \quad u_{\mathbf{k}} v_{\mathbf{k}}^* = g_{\mathbf{k}}^* = \frac{\Delta_{\mathbf{k}}^*}{2E_{\mathbf{k}}}, \quad u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$$

$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

$$E_{\text{BCS}} = \sum_{\mathbf{k}} \left( \xi_{\mathbf{k}} - \frac{\xi_{\mathbf{k}}^2}{E_{\mathbf{k}}} \right) - \sum_{\mathbf{k}} g_{\mathbf{k}}^* \Delta_{\mathbf{k}} = \sum_{\mathbf{k}} \left( \xi_{\mathbf{k}} - \frac{\xi_{\mathbf{k}}^2}{E_{\mathbf{k}}} \right) - \underbrace{2 \sum_{\mathbf{k}} g_{\mathbf{k}}^* \Delta_{\mathbf{k}}}_{-\sum_{\mathbf{k}} \Delta_{\mathbf{k}}^2 / 2E_{\mathbf{k}}} + \sum_{\mathbf{k}} g_{\mathbf{k}}^* \Delta_{\mathbf{k}}$$

$$E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2}$$

$$E_{\text{BCS}} = \sum_{\mathbf{k}} (\xi_{\mathbf{k}} - E_{\mathbf{k}}) + \sum_{\mathbf{k}} g_{\mathbf{k}}^* \Delta_{\mathbf{k}}$$

# 4.2.2 Ground State Energy

- for simplicity we assume for  $V_{\mathbf{k},\mathbf{k}'} = -V_0$  and  $\Delta_{\mathbf{k}} = \Delta$ )

$$E_{\text{BCS}} = \langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} - \mathcal{N}\mu | \Psi_{\text{BCS}} \rangle = \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} - E_{\mathbf{k}} + g_{\mathbf{k}}^* \Delta \}$$

- subtract mean energy of normal state at  $T = 0$  (making use of symmetry around  $\mu$ )

$$\langle \Psi_{\text{BCS}} | \mathcal{H}_n - \mathcal{N}\mu | \Psi_{\text{BCS}} \rangle = \lim_{\Delta \rightarrow 0} \langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} - \mathcal{N}\mu | \Psi_{\text{BCS}} \rangle = \sum_{\mathbf{k}} \xi_{\mathbf{k}} - |\xi_{\mathbf{k}}| = 2 \sum_{|\mathbf{k}| < \mathbf{k}_F} \xi_{\mathbf{k}}$$

$$\Delta E = \sum_{|\mathbf{k}| < \mathbf{k}_F} \xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta g_{\mathbf{k}}^* - 2 \xi_{\mathbf{k}} + \sum_{|\mathbf{k}| \geq \mathbf{k}_F} \xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta g_{\mathbf{k}}^*$$

- we use  $-\xi_{\mathbf{k}} = |\xi_{\mathbf{k}}|$  for  $|k| < k_F$  and  $E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}$

$$\Delta E = 2 \sum_{|\mathbf{k}| \geq \mathbf{k}_F} \left( \xi_{\mathbf{k}} - \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2} + \Delta g_{\mathbf{k}}^* \right) \stackrel{\Delta g_{\mathbf{k}}^* = \frac{\Delta^2}{2\sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}}}{=} 2 \sum_{|\mathbf{k}| \geq \mathbf{k}_F} \left( \xi_{\mathbf{k}} - \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2} + \frac{\Delta^2}{2\sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}} \right)$$

# 4.2.2 Ground State Energy

$$\Delta E = 2 \sum_{|\mathbf{k}| \geq k_F} \left( \xi_{\mathbf{k}} - \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2} + \frac{\Delta^2}{2\sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}} \right)$$

- replace summation by integration ..... after some algebra (see appendix H.3 in R. Gross, A. Marx, Festkörperphysik, 4. Auflage, de Gruyter (2022)):

$$\Delta E = E_{\text{cond}}(0) = -\frac{1}{4} D(E_F) \Delta^2(0)$$

$D(E_F)$  = DOS for both spin directions

interpretation of the result:

➤ number of Cooper pairs:

$$\frac{D(E_F)}{2} \Delta(0)$$

➤ average energy gain per Cooper pair:  $-\frac{\Delta(0)}{2}$

- compare to  $\mathcal{G}_s - \mathcal{G}_n = E_{\text{cond}}(0) = -B_{\text{cth}}^2(0)/2\mu_0$  (thermodynamics)



$$B_{\text{cth}}(0) = \sqrt{\frac{\mu_0 D(E_F) \Delta^2(0)}{2V}}$$

# 4.2.2 Ground State Energy

- condensation energy per volume:

$$\frac{E_{\text{cond}}(0)}{V} = -\frac{1}{4} \frac{D(E_F)}{V} \Delta^2(0) = -\frac{1}{4} N(E_F) \Delta^2(0)$$

with  $N(E_F) = \frac{3n}{2E_F}$  and  $\frac{\Delta(0)}{k_B T_c} = \frac{\pi}{e^\gamma} = 1.7638 \dots$  we obtain (result is derived later)

$$\frac{E_{\text{cond}}(0)}{V} = -\frac{3}{8} n \frac{\Delta^2(0)}{E_F} = \frac{3}{8} \left(\frac{\pi}{e^\gamma}\right)^2 \frac{(k_B T_c)^2}{E_F} = -1.167 n \frac{(k_B T_c)^2}{E_F}$$

→ average condensation energy per electron is of the order of  $(k_B T_c)^2 / E_F$

→ plausibility:

only a small fraction  $k_B T_c / E_F$  of the electrons is participating in pairing process and the average energy reduction per electron is about  $k_B T_c$

# Summary of Lecture No. 8 (1)

- BCS Hamilton operator:**

$$\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}$$

$$\hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} = n_{\mathbf{k}\sigma} = \text{particle number operator}$$

$$\varepsilon_{\mathbf{k}} = \xi_{\mathbf{k}} + \mu = \frac{\hbar^2 \mathbf{k}^2}{2m^*}$$

- Bardeen, Cooper and Schrieffer** used the following Ansatz for the ground state wave function (mean-field approach):

$$|\Psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}=\mathbf{k}_1, \dots, \mathbf{k}_M} (u_{\mathbf{k}} + v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger}) |0\rangle$$

$|u_{\mathbf{k}}|^2$ : probability that pair state ( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is empty


$|v_{\mathbf{k}}|^2$ : probability that pair state ( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is occupied

$$|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$$

coherent fermionic state

- expectation values:**  $\langle \Psi_{\text{BCS}} | \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} | \Psi_{\text{BCS}} \rangle = |v_{\mathbf{k}}|^2$        $\langle \Psi_{\text{BCS}} | \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} | \Psi_{\text{BCS}} \rangle = v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*$

$$\langle \Psi_{\text{BCS}} | \mathcal{H}_{\text{BCS}} | \Psi_{\text{BCS}} \rangle = \underbrace{2 \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} |v_{\mathbf{k}}|^2}_{= \bar{N} \varepsilon_{\mathbf{k}} \text{ (kinetic energy)}} + \underbrace{\sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}^* u_{\mathbf{k}'} u_{\mathbf{k}}^*}_{\text{interaction energy}}$$



determination of  $u_{\mathbf{k}}, v_{\mathbf{k}}$  by minimization of expectation value

# Summary of Lecture No. 8 (2)

- minimization of expectation value




$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right] \quad |v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$u_{\mathbf{k}} v_{\mathbf{k}}^* = g_{\mathbf{k}}^* = \frac{\Delta_{\mathbf{k}}^*}{2E_{\mathbf{k}}} \quad u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$$

probability that pair state  
( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is **empty/occupied**

pairing amplitude



$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'} = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

$$\Delta_{\mathbf{k}}^{\dagger} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

self-consistent gap equation

(solution simple for  $V_{\mathbf{k},\mathbf{k}'} = -V_0$ )

- gap equation for  $T = 0$

$$\Delta = \frac{\hbar\omega_D}{\sinh(2/V_0 D(E_F))} \approx 2\hbar\omega_D e^{-2/V_0 D(E_F)}$$

energy gap corresponds to binding energy estimated for single Cooper pair

$V_0 D(E_F) \ll 1$ : weak coupling approximation,  $\sinh x \approx \frac{1}{2} \exp x$

- ground state energy

$$E_{\text{cond}}(0) = -\frac{1}{4} D(E_F) \Delta^2(0)$$

$\left(\frac{1}{2} D(E_F) \Delta(0) \approx \text{number of pairs}, \Delta(0)/2 \approx \text{average binding energy per pair}\right)$





Walther  
Meißner  
Institut

**BAaW**

BAYERISCHE  
AKADEMIE  
DER  
WISSENSCHAFTEN

Technische  
Universität  
München

**TUM**

# Superconductivity and Low Temperature Physics I



**Lecture No. 9**

**R. Gross**

**© Walther-Meißner-Institut**

## 4. Microscopic Theory

### 4.1 Attractive Electron-Electron Interaction

#### 4.1.1 Phonon Mediated Interaction


#### 4.1.2 Cooper Pairs

#### 4.1.3 Symmetry of Pair Wavefunction

### 4.2 BCS Ground State

#### 4.2.1 The BCS Gap Equation

#### 4.2.2 Ground State Energy



#### 4.2.3 Bogoliubov-Valatin Transformation and Quasiparticle Excitations

### 4.3 Thermodynamic Quantities

### 4.4 Determination of the Energy Gap

#### 4.4.1 Specific Heat

#### 4.4.2 Tunneling Spectroscopy

### 4.5 Coherence Effects

# 4.2.3 The Bogoliubov-Valatin Transformation

- so far we have found the BCS ground state wave function, the energy gap, and the condensation energy at  $T = 0$
- **next step:**
  - determine the properties of the superconducting state **at finite temperature**
  - determine the **energy of excitations** out of the ground state
- **how to proceed?**
  - use BCS ground state as reference state
  - discuss effect of small deviations (e.g. by adding a small number of excitations to the ground state)
- we use the identities (with  $\delta g_{\mathbf{k}}, \delta g_{\mathbf{k}}^*$  being small)

$$\hat{c}_{-\mathbf{k}\downarrow}\hat{c}_{\mathbf{k}\uparrow} = \underbrace{\langle \hat{c}_{-\mathbf{k}\downarrow}\hat{c}_{\mathbf{k}\uparrow} \rangle}_{g_{\mathbf{k}}} + \underbrace{\hat{c}_{-\mathbf{k}\downarrow}\hat{c}_{\mathbf{k}\uparrow} - \langle \hat{c}_{-\mathbf{k}\downarrow}\hat{c}_{\mathbf{k}\uparrow} \rangle}_{\delta g_{\mathbf{k}}}$$

$$g_{\mathbf{k}} \equiv \langle c_{-\mathbf{k}\downarrow}c_{\mathbf{k}\uparrow} \rangle \neq 0$$

with pairing amplitude:

$$\hat{c}_{\mathbf{k}\uparrow}^{\dagger}\hat{c}_{-\mathbf{k}\downarrow}^{\dagger} = \underbrace{\langle \hat{c}_{\mathbf{k}\uparrow}^{\dagger}\hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \rangle}_{g_{\mathbf{k}}^*} + \underbrace{\hat{c}_{\mathbf{k}\uparrow}^{\dagger}\hat{c}_{-\mathbf{k}\downarrow}^{\dagger} - \langle \hat{c}_{\mathbf{k}\uparrow}^{\dagger}\hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \rangle}_{\delta g_{\mathbf{k}}^*}$$

$$g_{\mathbf{k}}^* \equiv \langle c_{-\mathbf{k}\downarrow}^{\dagger}c_{\mathbf{k}\uparrow}^{\dagger} \rangle \neq 0$$

as the particle number is usually very large, the fluctuations  $\delta g_{\mathbf{k}}, \delta g_{\mathbf{k}}^*$  are very small and we can neglect quadratic terms in  $\delta g_{\mathbf{k}}, \delta g_{\mathbf{k}}^*$

# 4.2.3 The Bogoliubov-Valatin Transformation

- rewriting of pair creation and annihilation operators in  $\mathcal{H}_{\text{BCS}} = \sum_{\sigma} \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'}^N V_{\mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow}$

$$\hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} = \underbrace{\langle \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \rangle}_{g_{\mathbf{k}}} + \underbrace{\hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} - \langle \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \rangle}_{\delta g_{\mathbf{k}}}$$

$$g_{\mathbf{k}} \equiv \langle c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} \rangle \neq 0$$

$$\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} = \underbrace{\langle \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \rangle}_{g_{\mathbf{k}}^*} + \underbrace{\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} - \langle \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \rangle}_{\delta g_{\mathbf{k}}^*}$$

$$g_{\mathbf{k}}^* \equiv \langle c_{-\mathbf{k}\downarrow}^{\dagger} c_{\mathbf{k}\uparrow}^{\dagger} \rangle \neq 0$$

pairing amplitude:

- insert into Hamiltonian and consider only terms linear in  $\delta g_{\mathbf{k}}^{(*)}$  (and after some math)

$$\mathcal{H}_{\text{BCS}} - N\mu = \sum_{\mathbf{k}, \sigma} \xi_{\mathbf{k}} n_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} \left[ g_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} + g_{\mathbf{k}'} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} - g_{\mathbf{k}}^* g_{\mathbf{k}'} \right]$$

- make use of pair potential  $\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} g_{\mathbf{k}'}$   $\Delta_{\mathbf{k}}^* \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} g_{\mathbf{k}'}^*$

$$\Rightarrow \mathcal{H}_{\text{BCS}} - N\mu = \sum_{\mathbf{k}, \sigma} \xi_{\mathbf{k}} n_{\mathbf{k}\sigma} - \sum_{\mathbf{k}} \left[ \Delta_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} + \Delta_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} - \Delta_{\mathbf{k}} g_{\mathbf{k}}^* \right]$$

# 4.2.3 The Bogoliubov-Valatin Transformation

- we use

$$\sum_{\mathbf{k}, \sigma} \xi_{\mathbf{k}} n_{\mathbf{k}\sigma} = \sum_{\mathbf{k}} \xi_{\mathbf{k}} (\hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} + \underbrace{\hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}}_{=1 - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^\dagger})$$

$$\sum_{\mathbf{k}, \sigma} \xi_{\mathbf{k}} n_{\mathbf{k}\sigma} = \sum_{\mathbf{k}} \xi_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} - \xi_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^\dagger + \xi_{\mathbf{k}}$$

$$\mathcal{H}_{\text{BCS}} - N\mu = \sum_{\mathbf{k}, \sigma} \xi_{\mathbf{k}} n_{\mathbf{k}\sigma} - \sum_{\mathbf{k}} [\Delta_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} + \Delta_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger - \Delta_{\mathbf{k}} g_{\mathbf{k}}^*]$$



$$\mathcal{H}_{\text{BCS}} - N\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + (\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow}) \begin{pmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}} \\ -\Delta_{\mathbf{k}}^* & -\xi_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix} \right\}$$

→ due to finite  $\Delta_{\mathbf{k}}, \Delta_{\mathbf{k}}^*$ , the Hamiltonian describes interacting electron gas with new quasiparticles consisting of *superposition of electron and hole states*

- derive excitation energies by *diagonalization of Hamiltonian*

→ *Bogoliubov-Valatin transformation*

→ define new fermionic operators  $\alpha_{\mathbf{k}}, \beta_{\mathbf{k}}^\dagger$  and  $\alpha_{\mathbf{k}}^\dagger, \beta_{\mathbf{k}}$  by unitary transformation (rotation)

# 4.2.3 The Bogoliubov-Valatin Transformation

- use unitarian matrix to rotate the energy matrix into eigenbasis of Bogoliubov quasiparticles

$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + \underbrace{(\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow})}_{\mathcal{C}_{\mathbf{k}}^\dagger} \underbrace{\begin{pmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}} \\ -\Delta_{\mathbf{k}}^* & -\xi_{\mathbf{k}} \end{pmatrix}}_{\mathcal{E}_{\mathbf{k}}} \underbrace{\begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix}}_{\mathcal{C}_{\mathbf{k}}} \right\} = \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + \mathcal{C}_{\mathbf{k}}^\dagger \mathcal{E}_{\mathbf{k}} \mathcal{C}_{\mathbf{k}} \}$$

spinors
energy matrix

$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + \underbrace{\mathcal{C}_{\mathbf{k}}^\dagger \mathcal{U}_{\mathbf{k}}}_{\mathcal{B}_{\mathbf{k}}^\dagger} \underbrace{\mathcal{U}_{\mathbf{k}}^\dagger \mathcal{E}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}}_{\tilde{\mathcal{E}}_{\mathbf{k}}} \underbrace{\mathcal{U}_{\mathbf{k}}^\dagger \mathcal{C}_{\mathbf{k}}}_{\mathcal{B}_{\mathbf{k}}} \right\} = \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + \mathcal{B}_{\mathbf{k}}^\dagger \tilde{\mathcal{E}}_{\mathbf{k}} \mathcal{B}_{\mathbf{k}} \} \quad \mathcal{U}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}^\dagger = 1$$

spinors of Bogoliubov quasiparticle operators:  $\mathcal{B}_{\mathbf{k}}^\dagger = (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) = \mathcal{C}_{\mathbf{k}}^\dagger \mathcal{U}_{\mathbf{k}} \quad \mathcal{B}_{\mathbf{k}} = (\alpha_{\mathbf{k}}, \beta_{-\mathbf{k}}^\dagger) = \mathcal{U}_{\mathbf{k}}^\dagger \mathcal{C}_{\mathbf{k}}$

appropriate unitary matrix to make transformed energy matrix  $\tilde{\mathcal{E}}_{\mathbf{k}} = \mathcal{U}_{\mathbf{k}}^\dagger \mathcal{E}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}$  diagonal:

$$\mathcal{U}_{\mathbf{k}} = \begin{pmatrix} u_{\mathbf{k}} & v_{\mathbf{k}}^* \\ -v_{\mathbf{k}} & u_{\mathbf{k}}^* \end{pmatrix} \quad \mathcal{U}_{\mathbf{k}}^\dagger = \begin{pmatrix} u_{\mathbf{k}}^* & -v_{\mathbf{k}}^* \\ v_{\mathbf{k}} & u_{\mathbf{k}} \end{pmatrix} \quad \Rightarrow \quad \mathcal{U}_{\mathbf{k}}^\dagger \mathcal{E}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}} = \tilde{\mathcal{E}}_{\mathbf{k}} = \begin{pmatrix} E_{\mathbf{k}} & D_{\mathbf{k}} \\ -D_{\mathbf{k}} & -E_{\mathbf{k}} \end{pmatrix} \quad \text{choose } u_{\mathbf{k}} \text{ and } v_{\mathbf{k}} \text{ such that off-diagonal terms vanish}$$

$$\Rightarrow \mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) \begin{pmatrix} E_{\mathbf{k}} & 0 \\ 0 & -E_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \alpha_{\mathbf{k}} \\ \beta_{-\mathbf{k}}^\dagger \end{pmatrix} \right\} \quad \text{eigenenergies } \pm E_{\mathbf{k}}$$

# 4.2.3 The Bogoliubov-Valatin Transformation

$$\begin{aligned}
 \mathcal{B}_{\mathbf{k}}^\dagger &= (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) = c_{\mathbf{k}}^\dagger u_{\mathbf{k}} = u_{\mathbf{k}}^T c_{\mathbf{k}}^\dagger & \Rightarrow & (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) = \begin{pmatrix} u_{\mathbf{k}} & -v_{\mathbf{k}} \\ v_{\mathbf{k}}^* & u_{\mathbf{k}}^* \end{pmatrix} \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow}^\dagger \\ \hat{c}_{-\mathbf{k}\downarrow} \end{pmatrix} \\
 \mathcal{B}_{\mathbf{k}} &= (\alpha_{\mathbf{k}}, \beta_{-\mathbf{k}}^\dagger) = u_{\mathbf{k}}^\dagger c_{\mathbf{k}} & \Rightarrow & (\alpha_{\mathbf{k}}, \beta_{-\mathbf{k}}^\dagger) = \begin{pmatrix} u_{\mathbf{k}}^* & -v_{\mathbf{k}}^* \\ v_{\mathbf{k}} & u_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \alpha_{\mathbf{k}} &= u_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow} - v_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\
 \beta_{-\mathbf{k}}^\dagger &= v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow} + u_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\
 \alpha_{\mathbf{k}}^\dagger &= u_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger - v_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow} \\
 \beta_{-\mathbf{k}} &= v_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow}^\dagger + u_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow}
 \end{aligned}$$

creation and annihilation operators for **Bogoliubov quasiparticles**: *symmetric and anti-symmetric superposition of electron and hole states with opposite momentum and spin*

- operators satisfy fermionic anti-commutation rules:  $\{\alpha_{\mathbf{k}}, \beta_{-\mathbf{k}'}^\dagger\} = \delta_{\mathbf{k}\mathbf{k}'}$  and  $\{\alpha_{\mathbf{k}}, \alpha_{\mathbf{k}'}\} = \{\alpha_{\mathbf{k}}^\dagger, \alpha_{\mathbf{k}'}^\dagger\} = 0$

- inverse transformation

$$\begin{aligned}
 \mathcal{B}_{\mathbf{k}}^\dagger u_{\mathbf{k}}^\dagger &= c_{\mathbf{k}}^\dagger u_{\mathbf{k}} u_{\mathbf{k}}^\dagger = c_{\mathbf{k}}^\dagger \Rightarrow c_{\mathbf{k}}^\dagger = (u_{\mathbf{k}}^\dagger)^T \mathcal{B}_{\mathbf{k}}^\dagger & \Rightarrow & (\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow}) = \begin{pmatrix} u_{\mathbf{k}}^* & v_{\mathbf{k}} \\ -v_{\mathbf{k}}^* & u_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \alpha_{\mathbf{k}}^\dagger \\ \beta_{-\mathbf{k}} \end{pmatrix} \\
 u_{\mathbf{k}} \mathcal{B}_{\mathbf{k}} &= u_{\mathbf{k}} u_{\mathbf{k}}^\dagger c_{\mathbf{k}} = c_{\mathbf{k}} \Rightarrow c_{\mathbf{k}} = u_{\mathbf{k}} \mathcal{B}_{\mathbf{k}} & \Rightarrow & (\hat{c}_{\mathbf{k}\uparrow}, \hat{c}_{-\mathbf{k}\downarrow}^\dagger) = \begin{pmatrix} u_{\mathbf{k}} & v_{\mathbf{k}}^* \\ -v_{\mathbf{k}} & u_{\mathbf{k}}^* \end{pmatrix} \begin{pmatrix} \alpha_{\mathbf{k}} \\ \beta_{-\mathbf{k}}^\dagger \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \hat{c}_{\mathbf{k}\uparrow}^\dagger &= u_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + v_{\mathbf{k}} \beta_{-\mathbf{k}} \\
 \hat{c}_{-\mathbf{k}\downarrow} &= -v_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + u_{\mathbf{k}} \beta_{-\mathbf{k}} \\
 \hat{c}_{\mathbf{k}\uparrow} &= u_{\mathbf{k}} \alpha_{\mathbf{k}} + v_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger \\
 \hat{c}_{-\mathbf{k}\downarrow}^\dagger &= -v_{\mathbf{k}} \alpha_{\mathbf{k}} + u_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger
 \end{aligned}$$


# 4.2.3 The Bogoliubov-Valatin Transformation

## Bogoliubov quasiparticles

$$\begin{aligned} \alpha_{\mathbf{k}} &= u_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow} - v_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\ \beta_{-\mathbf{k}}^\dagger &= v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow} + u_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\ \alpha_{\mathbf{k}}^\dagger &= u_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger - v_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow} \\ \beta_{-\mathbf{k}} &= v_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow}^\dagger + u_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow} \end{aligned}$$


$$\begin{aligned} \xi_{\mathbf{k}} &= 0 \\ |u_{\mathbf{k}}|^2 &= |v_{\mathbf{k}}|^2 = \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \alpha_{\mathbf{k}} &= 1/\sqrt{2}(\hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow}^\dagger) \\ \beta_{-\mathbf{k}}^\dagger &= 1/\sqrt{2}(\hat{c}_{\mathbf{k}\uparrow} + \hat{c}_{-\mathbf{k}\downarrow}^\dagger) \\ \alpha_{\mathbf{k}}^\dagger &= 1/\sqrt{2}(\hat{c}_{\mathbf{k}\uparrow}^\dagger - \hat{c}_{-\mathbf{k}\downarrow}) \\ \beta_{-\mathbf{k}} &= 1/\sqrt{2}(\hat{c}_{\mathbf{k}\uparrow}^\dagger + \hat{c}_{-\mathbf{k}\downarrow}) \end{aligned}$$




→ symmetric and anti-symmetric superposition of electron and hole states with opposite spin direction

→  $|u_{\mathbf{k}}|^2 =$  hole fraction,  $|v_{\mathbf{k}}|^2 =$  electron fraction



→ *reduces* the total momentum by  $\mathbf{k}$  and the total spin by  $\hbar/2$   
*hole-like excitation*



→ *increases* the total momentum by  $\mathbf{k}$  and the total spin by  $\hbar/2$   
*particle-like excitation*



# 4.2.3 The Bogoliubov-Valatin Transformation

determine  $|u_{\mathbf{k}}|^2$  and  $|v_{\mathbf{k}}|^2$  by Bogoliubov-Valatin transformation

BCS Hamiltonian

$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + (\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow}) \begin{pmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}} \\ -\Delta_{\mathbf{k}}^* & -\xi_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix} \right\}$$

$$\begin{aligned} \hat{c}_{\mathbf{k}\uparrow}^\dagger &= u_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + v_{\mathbf{k}} \beta_{-\mathbf{k}} \\ \hat{c}_{-\mathbf{k}\downarrow} &= -v_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + u_{\mathbf{k}} \beta_{-\mathbf{k}} \\ \hat{c}_{\mathbf{k}\uparrow} &= u_{\mathbf{k}} \alpha_{\mathbf{k}} + v_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger &= -v_{\mathbf{k}} \alpha_{\mathbf{k}} + u_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger \end{aligned}$$

- replace operators by Bogoliubov quasiparticle operators  $\rightarrow$  resulting Hamiltonian:

$$\begin{aligned} \mathcal{H}_{\text{BCS}} - \mathcal{N}\mu &= \sum_{\mathbf{k}} [2\xi_{\mathbf{k}} v_{\mathbf{k}}^2 - \Delta_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}}^* - \Delta_{\mathbf{k}}^\dagger u_{\mathbf{k}}^* v_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^\dagger] \\ &+ \sum_{\mathbf{k}} [\xi_{\mathbf{k}} (u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2) + \Delta_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}}^* + \Delta_{\mathbf{k}}^\dagger u_{\mathbf{k}}^* v_{\mathbf{k}}] \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} \\ &+ \sum_{\mathbf{k}} [\xi_{\mathbf{k}} (u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2) + \Delta_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}}^* + \Delta_{\mathbf{k}}^\dagger u_{\mathbf{k}}^* v_{\mathbf{k}}] \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}} \\ &+ \sum_{\mathbf{k}} \left[ 2\xi_{\mathbf{k}} u_{\mathbf{k}}^* v_{\mathbf{k}}^* + \Delta_{\mathbf{k}} v_{\mathbf{k}}^{*2} - \Delta_{\mathbf{k}}^\dagger u_{\mathbf{k}}^{*2} \right] \beta_{-\mathbf{k}} \alpha_{\mathbf{k}} \\ &+ \sum_{\mathbf{k}} \left[ 2\xi_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}} + \Delta_{\mathbf{k}}^\dagger v_{\mathbf{k}}^2 - \Delta_{\mathbf{k}} u_{\mathbf{k}}^2 \right] \alpha_{\mathbf{k}}^\dagger \beta_{-\mathbf{k}}^\dagger \end{aligned} \quad \left. \vphantom{\sum_{\mathbf{k}}} \right\} [\dots] = \mathbf{0}!$$

- $\triangleright$  we have to set expressions marked in red to zero to keep only diagonal terms
- $\triangleright$   $\alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}}$  and  $\beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}}$  = quasiparticle number operators

# 4.2.3 The Bogoliubov-Valatin Transformation



$$2\xi_{\mathbf{k}}u_{\mathbf{k}}v_{\mathbf{k}} + \Delta_{\mathbf{k}}^*v_{\mathbf{k}}^2 - \Delta_{\mathbf{k}}u_{\mathbf{k}}^2 = 0 \quad \text{and} \quad 2\xi_{\mathbf{k}}u_{\mathbf{k}}^*v_{\mathbf{k}}^* + \Delta_{\mathbf{k}}v_{\mathbf{k}}^{*2} - \Delta_{\mathbf{k}}^*u_{\mathbf{k}}^{*2} = 0$$

- multiply by  $\Delta_{\mathbf{k}}^*/u_{\mathbf{k}}^2$  ( $\Delta_{\mathbf{k}}/u_{\mathbf{k}}^{*2}$ ), solve the resulting quadratic eqn. for  $\Delta_{\mathbf{k}}^*v_{\mathbf{k}}/u_{\mathbf{k}}$  ( $\Delta_{\mathbf{k}}v_{\mathbf{k}}^*/u_{\mathbf{k}}^*$ )

$$2\xi_{\mathbf{k}}u_{\mathbf{k}}v_{\mathbf{k}}\frac{\Delta_{\mathbf{k}}^*}{u_{\mathbf{k}}^2} + \Delta_{\mathbf{k}}^*v_{\mathbf{k}}^2\frac{\Delta_{\mathbf{k}}^*}{u_{\mathbf{k}}^2} - \Delta_{\mathbf{k}}u_{\mathbf{k}}^2\frac{\Delta_{\mathbf{k}}^*}{u_{\mathbf{k}}^2} = \left(\Delta_{\mathbf{k}}^*\frac{v_{\mathbf{k}}}{u_{\mathbf{k}}}\right)^2 + 2\xi_{\mathbf{k}}\left(\Delta_{\mathbf{k}}^*\frac{v_{\mathbf{k}}}{u_{\mathbf{k}}}\right) + |\Delta_{\mathbf{k}}|^2 = 0$$



$$\left(\Delta_{\mathbf{k}}^*\frac{v_{\mathbf{k}}}{u_{\mathbf{k}}}\right)_{1,2} = \left(\Delta_{\mathbf{k}}\frac{v_{\mathbf{k}}^*}{u_{\mathbf{k}}^*}\right)_{1,2} = -\xi_{\mathbf{k}} \pm \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2} = -\xi_{\mathbf{k}} + E_{\mathbf{k}} \quad \text{with} \quad E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

negative sign is unphysical

→ corresponds to solution with maximum energy

note that the phases of  $u_{\mathbf{k}}$ ,  $v_{\mathbf{k}}$  and  $\Delta_{\mathbf{k}}^*$  ( $u_{\mathbf{k}}^*$ ,  $v_{\mathbf{k}}^*$  and  $\Delta_{\mathbf{k}}$ ), although arbitrary, are related, since the quantity on the r.h.s. is real

→ the relative phase of  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  must be fixed and must be the phase of  $\Delta_{\mathbf{k}}^*$

→ we can choose  $u_{\mathbf{k}}$  real and use  $v_{\mathbf{k}} = |v_{\mathbf{k}}|e^{i\varphi}$ , the phase of  $v_{\mathbf{k}}$  corresponds to that of  $\Delta_{\mathbf{k}}^*$



$$\left|\frac{v_{\mathbf{k}}}{u_{\mathbf{k}}}\right| = \frac{E_{\mathbf{k}} - \xi_{\mathbf{k}}}{|\Delta_{\mathbf{k}}|}$$

# 4.2.3 The Bogoliubov-Valatin Transformation

- with the condition  $\left| \frac{v_{\mathbf{k}}}{u_{\mathbf{k}}} \right| = \frac{E_{\mathbf{k}} - \xi_{\mathbf{k}}}{|\Delta_{\mathbf{k}}|}$  and the normalization condition  $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$  we obtain

$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right] \quad |v_{\mathbf{k}}|^2 = \frac{1}{2} \left[ 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right]$$

$$u_{\mathbf{k}} v_{\mathbf{k}}^* = g_{\mathbf{k}}^* = \frac{\Delta_{\mathbf{k}}^*}{2E_{\mathbf{k}}} \quad u_{\mathbf{k}}^* v_{\mathbf{k}} = g_{\mathbf{k}} = \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}}$$

probability that pair state ( $\mathbf{k} \uparrow, -\mathbf{k} \downarrow$ ) is empty/occupied

pairing amplitude

$$E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$


$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'} = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

$$\Delta_{\mathbf{k}}^* \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}^*}{2E_{\mathbf{k}'}}$$

self-consistent gap equation

# 4.2.3 The Bogoliubov-Valatin Transformation

excitation spectrum of Bogoliubov quasiparticles and energy gap




**quasiparticle excitations: superposition of electron and hole states**

**reason:** single particle excitation with wave vector  $\mathbf{k}$  can only exist if at the same time, if there is a hole with wave vector  $-\mathbf{k}$ , otherwise there would be a pair state


excitation energy

$$E_{\mathbf{k}} = \pm \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$



$$\begin{aligned} \alpha_{\mathbf{k}} &= u_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow} - v_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\ \beta_{-\mathbf{k}}^\dagger &= v_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow} + u_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\ \alpha_{\mathbf{k}}^\dagger &= u_{\mathbf{k}} \hat{c}_{\mathbf{k}\uparrow}^\dagger - v_{\mathbf{k}} \hat{c}_{-\mathbf{k}\downarrow} \\ \beta_{-\mathbf{k}} &= v_{\mathbf{k}}^* \hat{c}_{\mathbf{k}\uparrow}^\dagger + u_{\mathbf{k}}^* \hat{c}_{-\mathbf{k}\downarrow} \end{aligned}$$

# 4.2.3 The Bogoliubov-Valatin Transformation



# 4.2.3 The Bogoliubov-Valatin Transformation

## reformulation of the BCS Hamilton operator

• we start from the Hamiltonian  $\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) \begin{pmatrix} E_{\mathbf{k}} & 0 \\ 0 & -E_{\mathbf{k}} \end{pmatrix} \begin{pmatrix} \alpha_{\mathbf{k}} \\ \beta_{-\mathbf{k}}^\dagger \end{pmatrix} \right\}$

$$\begin{aligned} \mathcal{H}_{\text{BCS}} - \mathcal{N}\mu &= \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* \} + \sum_{\mathbf{k}} \left\{ E_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} - E_{\mathbf{k}} \underbrace{\beta_{-\mathbf{k}} \beta_{-\mathbf{k}}^\dagger}_{=1 - \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}}} \right\} \\ &= \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* \} + \sum_{\mathbf{k}} \{ E_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} - E_{\mathbf{k}} + E_{\mathbf{k}} \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}} \} \end{aligned}$$



$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \underbrace{\sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* \}}_{\text{mean-field contribution}} + \underbrace{\sum_{\mathbf{k}} \{ E_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + E_{\mathbf{k}} \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}} \}}_{\text{spinless Fermion system}}$$

→ minimization of free energy yields BCS gap equation for finite  $T$

**mean-field contribution ( $T = 0$ )**

differs from the normal state value by the condensation energy

contribution of **spinless Fermion system** with two kind of **quasiparticles** described by operators  $\alpha_{\mathbf{k}}^\dagger, \alpha_{\mathbf{k}}$  and  $\beta_{-\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}$  and excitation energies  $\pm E_{\mathbf{k}}$

spinless **quasiparticles** since they consist of superposition of spin- $\uparrow$  and spin- $\downarrow$  electrons

## 4.2.3 The Bogoliubov-Valatin Transformation

- note that the **Bogoliubov quasiparticles** are not part of the BCS ground state, as is evident from

$$\alpha_{\mathbf{k}}|\Psi_{\text{BCS}}\rangle = 0$$

$$\beta_{-\mathbf{k}}|\Psi_{\text{BCS}}\rangle = 0$$

- the occupation probability of the Bogoliubov particles is given by the Fermi-Dirac distribution

$$\langle \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} \rangle = \langle \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}} \rangle = f(E_{\mathbf{k}}) = \frac{1}{\exp(E_{\mathbf{k}}/k_{\text{B}}T) + 1}$$

# 4.2.3 The BCS Gap Equation and QP Excitations

determination of temperature dependence of  $\Delta$  by minimization of free energy

- Hamiltonian has two terms:  $\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \underbrace{\sum_{\mathbf{k}} \{\xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^*\}}_{\text{constant term } \mathcal{H}_0} + \underbrace{\sum_{\mathbf{k}} \{E_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + E_{\mathbf{k}} \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}}\}}_{\text{term of free Fermi gas composed of two kind of fermions with energy } E_{\mathbf{k}}}$

- grand canonical partition function:

$$Z = e^{-\mathcal{H}_0/k_B T} \prod_{\mathbf{k}} (1 + e^{-E_{\mathbf{k}}/k_B T})(1 + e^{E_{\mathbf{k}}/k_B T}) = e^{-\mathcal{F}/Nk_B T}$$

(since  $\mathcal{F} = -Nk_B T \ln Z$ )

partition function of an ideal Fermi gas:

$$Z = \prod_{\mathbf{k}} \sum_{n_{\mathbf{k}}=0,1} \exp(-n_{\mathbf{k}}(\epsilon_{\mathbf{k}} - \mu)/k_B T)$$

$$= \prod_{\mathbf{k}} [1 + \exp(-(\epsilon_{\mathbf{k}} - \mu)/k_B T)]$$

- solve for free energy  $\mathcal{F}$  :

$$\frac{\mathcal{F}}{N} = \mathcal{H}_0 - k_B T \sum_{\mathbf{k}} [\ln(1 + e^{-E_{\mathbf{k}}/k_B T}) + \ln(1 + e^{E_{\mathbf{k}}/k_B T})]$$

- minimize free energy regarding variation of  $\Delta_{\mathbf{k}}$ :

$$\frac{\partial \mathcal{F}}{\partial \Delta_{\mathbf{k}}} = 0, \quad \frac{\partial \mathcal{F}}{\partial \Delta_{\mathbf{k}}^\dagger} = 0$$



# 4.2.3 The BCS Gap Equation and QP Excitations

$$\frac{\partial(\mathcal{F}/N)}{\partial\Delta_{\mathbf{k}}} = 0 = \frac{\partial}{\partial\Delta_{\mathbf{k}}} \left\{ \mathcal{H}_0 - k_B T \sum_{\mathbf{k}} [\ln(1 + e^{-E_{\mathbf{k}}/k_B T}) + \ln(1 + e^{E_{\mathbf{k}}/k_B T})] \right\}$$

$$\mathcal{H}_0 = \xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^*$$

$$\Rightarrow g_{\mathbf{k}}^* + \underbrace{\frac{\partial E_{\mathbf{k}}}{\partial\Delta_{\mathbf{k}}}}_{=\Delta_{\mathbf{k}}^*/2E_{\mathbf{k}}} \underbrace{\left[ \frac{e^{-E_{\mathbf{k}}/k_B T}}{1 + e^{-E_{\mathbf{k}}/k_B T}} - \frac{e^{E_{\mathbf{k}}/k_B T}}{1 + e^{E_{\mathbf{k}}/k_B T}} \right]}_{=-\tanh(E_{\mathbf{k}}/2k_B T)} = 0$$

$$\Delta_{\mathbf{k}} \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}}$$

$$g_{\mathbf{k}}^* = \frac{\Delta_{\mathbf{k}}^*}{2E_{\mathbf{k}}} \tanh\left(\frac{E_{\mathbf{k}}}{2k_B T}\right) = u_{\mathbf{k}} v_{\mathbf{k}}^* \tanh\left(\frac{E_{\mathbf{k}}}{2k_B T}\right)$$

$$u_{\mathbf{k}} v_{\mathbf{k}}^* = \frac{\Delta_{\mathbf{k}}^*}{2E_{\mathbf{k}}}$$

**pairing susceptibility/amplitude: ability of the electron system to form pairs**

- we use  $\Delta_{\mathbf{k}}^* \equiv - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} g_{\mathbf{k}'}^*$  and obtain:

$$\Delta_{\mathbf{k}}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}} \tanh\left(\frac{E_{\mathbf{k}'}}{2k_B T}\right)$$

**BCS gap equation**

- set of equations for variables  $\Delta_{\mathbf{k}}$
- equations are nonlinear, since  $E_{\mathbf{k}}$  depends on  $\Delta_{\mathbf{k}}$
- solve numerically, analytical solutions in limiting cases

# 4.2.3 The BCS Gap Equation and QP Excitations

*energy gap  $\Delta$  and transition temperature  $T_c$*

- trivial solution:  $\Delta_{\mathbf{k}} = 0$ , results in  $v_{\mathbf{k}} = 1$  for  $\xi_{\mathbf{k}} < 0$  and  $v_{\mathbf{k}} = 0$  for  $\xi_{\mathbf{k}} > 0$

→ intuitive expectation for normal state

- non-trivial solution: we use approximations  $V_{\mathbf{k},\mathbf{k}'} = -V_0$  and  $\Delta_{\mathbf{k}} = \Delta$

$$\Delta_{\mathbf{k}}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}^*}{2E_{\mathbf{k}'}} \tanh \left( \frac{E_{\mathbf{k}'}}{2k_B T} \right) \quad \Rightarrow \quad 1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2E_{\mathbf{k}'}} \tanh \left( \frac{E_{\mathbf{k}'}}{2k_B T} \right)$$

- we use pair density of states  $\tilde{D}(E) = D(E)/2$  and change from summation to integration

**simple solutions for**

- (i)  $T \rightarrow 0$**
- (ii)  $T \rightarrow T_c$**

# 4.2.3 The BCS Gap Equation and QP Excitations

i. **solution for  $T \rightarrow 0$ :** (already discussed above for  $V_{\mathbf{k},\mathbf{k}'} = -V_0$  and  $\Delta_{\mathbf{k}} = \Delta$ )

$$1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2E_{\mathbf{k}'}} \underbrace{\tanh\left(\frac{E_{\mathbf{k}'}}{2k_B T}\right)}_{=1 \text{ for } T \rightarrow 0} \quad \text{transforming sum into integration} \quad 1 = \frac{V_0 D(E_F)}{2} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{d\xi}{2\sqrt{\xi_{\mathbf{k}}^2 + |\Delta(0)|^2}} \quad \text{with } \tilde{D}(E) \simeq D(E_F)/2$$

$$1 = \frac{V_0 D(E_F)}{4} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{d\xi}{\sqrt{\xi_{\mathbf{k}}^2 + |\Delta(0)|^2}} = \frac{V_0 D(E_F)}{4} \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta(0)}\right) \Big|_{-\hbar\omega_D}^{+\hbar\omega_D} = \frac{V_0 D(E_F)}{2} \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta(0)}\right)$$

• solve for  $\Delta$ :

$$\Rightarrow \Delta(0) = \frac{\hbar\omega_D}{\sinh(2/V_0 D(E_F))} \simeq 2\hbar\omega_D e^{-2/V_0 D(E_F)}$$

$V_0 D(E_F) \ll 1$ : weak coupling approximation

• compare to expression derived for energy of two interacting electrons (“Gedanken” experiment):

$$E \simeq 2E_F - 2\hbar\omega_D e^{-4/V_0 D(E_F)}$$

factor 2 in argument of exponential function since we have assumed that the two additional electrons are in interval between  $E_F$  and  $E_F + \hbar\omega_D$  and not between  $E_F - \hbar\omega_D$  and  $E_F + \hbar\omega_D$


# 4.2.3 The BCS Gap Equation and QP Excitations

ii. solution for  $T \rightarrow T_c$ :  $E_{\mathbf{k}} \rightarrow |\xi_{\mathbf{k}}|$  since  $\Delta_{\mathbf{k}} \rightarrow 0$

$$1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2E_{\mathbf{k}'}} \tanh\left(\frac{E_{\mathbf{k}'}}{2k_B T}\right) \Rightarrow 1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2\xi_{\mathbf{k}'}} \tanh\left(\frac{\xi_{\mathbf{k}'}}{2k_B T}\right)$$

$$1 = \frac{V_0 D(E_F)}{4} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{1}{\xi_{\mathbf{k}'}} \tanh\left(\frac{\xi_{\mathbf{k}'}}{2k_B T_c}\right) d\xi = \frac{V_0 D(E_F)}{4} \int_{-\hbar\omega_D/2k_B T_c}^{\hbar\omega_D/2k_B T_c} \frac{\tanh x}{x} dx \quad \text{with } x = \xi_{\mathbf{k}}/2k_B T_c$$

- integral gives  $2 \ln(p \hbar\omega_D/2k_B T_c)$  with  $p = \frac{2e^\gamma}{\pi} \simeq 1.13$  and  $\gamma = 0.577 \dots$  (Euler constant)

  $k_B T_c = 1.13 \hbar\omega_D e^{-2/V_0 D(E_F)}$

critical temperature is proportional to Debye frequency  $\omega_D \propto 1/\sqrt{M}$

**→ explains isotope effect !!**

# 4.2.3 The BCS Gap Equation and QP Excitations

relation between energy gap at zero temperature and critical temperature

$$\Delta(0) \simeq 2\hbar\omega_D e^{-2/V_0D(E_F)} \longleftrightarrow k_B T_c = \frac{2e^\gamma}{\pi} \hbar\omega_D e^{-2/V_0D(E_F)}$$

$$\frac{\Delta(0)}{k_B T_c} = \frac{\pi}{e^\gamma} = 1.764$$

key prediction of BCS theory

	$T_c$ (K)	$2\Delta(0)$ (meV)	$2\Delta(0)/k_B T_c$		$T_c$ (K)	$2\Delta(0)$ (meV)	$2\Delta(0)/k_B T_c$
Al	1.19	0.36	$3.5 \pm 0.1$	In	3.4	1.05	$3.5 \pm 0.1$
Nb	9.2	2.90	3.6	Hg	4.15	1.65	$4.6 \pm 0.1$
Pb	7.2	2.70	$4.3 \pm 0.05$	Sn	3.72	1.15	$3.5 \pm 0.1$
Ta	4.29	1.30	$3.5 \pm 0.1$	Tl	2.39	0.75	$3.6 \pm 0.1$
NbN	15	4.65	3.6	Nb <sub>3</sub> Sn	18	6.55	4.2
NbSe <sub>2</sub>	7	2.2	3.7	MgB <sub>2</sub>	40	3.6-15	1.1 – 4.5

considerable deviations for so-called „strong-coupling“ superconductors:


→  $V_0D(E_F) \ll 1$  is no longer a good approximation

# 4.2.3 The BCS Gap Equation and QP Excitations

solution for  $0 < T < T_c$  (numerical solution of integral)

$$1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2E_{\mathbf{k}'}} \tanh \left( \frac{E_{\mathbf{k}'}}{2k_B T} \right) \quad \text{sum} \Rightarrow \text{integral}$$

$$1 = \frac{V_0 D(E_F)}{2} \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{1}{2E_{\mathbf{k}}} \tanh \left( \frac{E_{\mathbf{k}}}{2k_B T} \right) d\xi_{\mathbf{k}}$$



good approximation close to  $T_c$ :

$$\frac{\Delta(T)}{\Delta(0)} \simeq 1.74 \left( 1 - \frac{T}{T_c} \right)^{1/2}$$

(characteristic result of mean-field theory)

B. Mühlischlegel, *Die thermodynamischen Funktionen des Supraleiters*, Z. Phys. 115, 313–327 (1959).

# 4.2.3 The BCS Gap Equation and QP Excitations

## strong electron-phonon coupling

- BCS results are valid only for weak coupling:  $V_0 D(E_F) \ll 1$
- for  $V_0 D(E_F) \gtrsim 0.2$  a more elaborate treatment is required

**phonons have influence on electrons but also electrons change e.g. phonon frequencies**

### • Eliashberg theory

- replace coupling constant  $\lambda = V_0 D(E_F)$  by

$$\lambda(\omega) = 2 \int_0^{\infty} \frac{\alpha^2(\omega) F(\omega)}{\omega} d\omega$$

$F(\omega)$ : phonon density of states

$\alpha(\omega)$ : matrix element of the electron-phonon interaction

G. M. Eliashberg, *Interactions Between Electrons and Lattice Vibrations in a Superconductor*, Zh. Éksp. Teor. Fiz. 38, 966 (1960) [Sov. Phys. JETP 11, 696-702 (1960)].

### • McMillan approximation

- several attempts have been made to improve prediction for  $T_c$  using strong coupling theory, e.g. by McMillan:

$$T_c = \frac{\hbar\omega_D}{1.45} \exp\left(\frac{1.04(1 + \lambda)}{\lambda - \mu^*(1 + 0.62 \lambda)}\right)$$


$\mu^*$ : matrix element of the short-range screened Coulomb repulsion

W. L. McMillan, *Transition Temperature of Strong-Coupled Superconductors*, Phys. Rev. 167, 331 (1968).

# 4.2.3 Energy Gap and Excitation Spectrum

dispersion of excitations (Bogoliubov quasiparticles) from the superconducting ground state

→ *excitations represent superpositions of electron- and hole-type single particle states*  
 (reason: single particle excitation with  $\mathbf{k}$  can only exist if there is hole with  $-\mathbf{k}$ , if not, Cooper pair would form)



excitation energy

$$E_{\mathbf{k}} = E_{-\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

- break up of Cooper pair requires energy  $2E_{\mathbf{k}}$
- $\Delta$  represents energy gap for quasiparticle excitation from ground state  
 → minimum excitation energy

equal superposition of electron with wave vector  $\mathbf{k}$  and hole with wave vector  $-\mathbf{k}$




# 4.2.3 Energy Gap and Excitation Spectrum

## density of states

- conservation of states on transition to sc state requires  $D_s(E_{\mathbf{k}})dE_{\mathbf{k}} = D_n(\xi_{\mathbf{k}})d\xi_{\mathbf{k}}$
- close to  $E_F$ :  $D_n(\xi_{\mathbf{k}}) \simeq D_n(E_F) = \text{const.}$

$$D_s(E_{\mathbf{k}}) = D_n(\xi_{\mathbf{k}}) \frac{d\xi_{\mathbf{k}}}{dE_{\mathbf{k}}} = \begin{cases} D_n(E_F) \frac{E_{\mathbf{k}}}{\sqrt{E_{\mathbf{k}}^2 - \Delta^2}} & \text{for } E_{\mathbf{k}} > \Delta \\ 0 & \text{for } E_{\mathbf{k}} < \Delta \end{cases}$$



- measurement by tunneling spectroscopy

I. Giaever,  
Phys. Rev. **126**, 941 (1962)

# 4.3 Thermodynamic Quantities

- occupation probability of qp-excitations is given by  $f(E_{\mathbf{k}}) = [\exp(E_{\mathbf{k}}/k_B T) + 1]^{-1}$

→ i.e. by  $\Delta_{\mathbf{k}}(T)$ , which is contained in  $E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2(T)}$

- **entropy of electronic system** (determined only by the occupation probability → is fixed by  $\Delta_{\mathbf{k}}$ )

$$S_s = -2k_B \sum_{\mathbf{k}} \left\{ \underbrace{[1 - f(E_{\mathbf{k}})] \ln[1 - f(E_{\mathbf{k}})]}_{\text{hole like}} + \underbrace{f(E_{\mathbf{k}}) \ln[f(E_{\mathbf{k}})]}_{\text{electron like}} \right\}$$

$$S = -k_B \sum_n p_n \ln p_n$$

- **heat capacity:**  $C_s = T \left( \frac{\partial S_s}{\partial T} \right)_{p.B}$

after some math:

$$C_s = \frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \left( \underbrace{E_{\mathbf{k}}^2}_{\text{results from redistribution of qp on available energy levels}} - \underbrace{\frac{1}{2} T \frac{d\Delta_{\mathbf{k}}^2(T)}{dT}}_{\text{results from } T\text{-dependence of energy gap}} \right)$$

**Yosida function:**

$$Y(T) = \frac{1}{D(E_F)} \sum_{\mathbf{k}} -\frac{\partial f(E_{\mathbf{k}}, T)}{\partial E_{\mathbf{k}}} = \frac{1}{4k_B T} \int_{-\mu}^{\infty} \frac{d\xi_{\mathbf{k}}}{\cosh^2(\xi_{\mathbf{k}}/2k_B T)}$$

appears in many thermodynamic properties

$Y(T)$  describes the  $T$ -dependence of the qp excitations (normal fluid density):  $n_n(T) = n Y(T)$

# 4.3 Thermodynamic Quantities

## discussion of limiting cases

### i. $T \ll T_c$ :

- since  $\Delta_{\mathbf{k}}(T) \simeq \Delta_{\mathbf{k}}(0) \gg k_B T$ , there are only a few thermally excited qp
- we use approximations  $d\Delta_{\mathbf{k}}^2(T)/dT \simeq 0$  and  $f(E_{\mathbf{k}}) = [\exp(E_{\mathbf{k}}/k_B T) + 1]^{-1} \simeq \exp(-E_{\mathbf{k}}/k_B T)$
- we assume  $\Delta_{\mathbf{k}} = \Delta$  for simplicity and transfer sum into an integral  
(we use  $\Delta^2 + \xi_{\mathbf{k}}^2 = \Delta^2(1 + \xi_{\mathbf{k}}^2/\Delta^2) \simeq \Delta^2$  and  $\sqrt{\Delta^2 + \xi_{\mathbf{k}}^2} = \Delta\sqrt{1 + \xi_{\mathbf{k}}^2/\Delta^2} \simeq \Delta + \xi_{\mathbf{k}}^2/2\Delta$ , as  $\partial f(E_{\mathbf{k}})/\partial E_{\mathbf{k}}$  has significant weight only for small values of  $\xi_{\mathbf{k}}/\Delta$ )

$$C_s = \frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \left( E_{\mathbf{k}}^2 - \frac{1}{2} T \frac{d\Delta_{\mathbf{k}}^2(T)}{dT} \right)$$

$$E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2(T)}$$

$$C_s = \frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \left( E_{\mathbf{k}}^2 - \frac{1}{2} T \frac{d\Delta^2(T)}{dT} \right) \simeq \frac{D(E_F)}{k_B T^2} \Delta^2(0) \int_0^{\infty} e^{-\sqrt{\Delta^2 + \xi_{\mathbf{k}}^2}/k_B T} d\xi_{\mathbf{k}}$$

$$\frac{\Delta(0)}{k_B T_c} = 1.76$$

$$C_s \simeq \frac{D(E_F)}{k_B T^2} \Delta^2(0) e^{-\Delta(0)/k_B T} \underbrace{\int_0^{\infty} e^{-\xi_{\mathbf{k}}^2/2\Delta(0)k_B T} d\xi_{\mathbf{k}}}_{\sqrt{\pi k_B T \Delta(0)}/2}$$

$$C_s \propto T^{-\frac{3}{2}} e^{-\frac{\Delta(0)}{k_B T}} \propto T^{-\frac{3}{2}} e^{-1.76 \frac{T_c}{T}} @ T \ll T_c$$


**exponential decrease of heat capacity at low T**

# 4.3 Thermodynamic Quantities

specific heat of superconductors at  $T \ll T_c$ :

exponential decrease of  $C_s$  with decreasing  $T$ :

$$C_s \propto T^{-\frac{3}{2}} e^{-\frac{\Delta(0)}{k_B T}} \propto T^{-\frac{3}{2}} e^{-1.76 \frac{T_c}{T}} \quad @ T \ll T_c$$





M. A. Biondi, A. T. Forrester, M. P. Garfunkel, C. B. Satterthwaite, Rev. Mod. Phys. **30**, 1109-1136 (1958)

# 4.3 Thermodynamic Quantities

ii.  $0.5 < T/T_c < 1$ :

$\Delta(T)$  decreases with increasing  $T$   $\rightarrow$  there is a rapid increase of the number of thermally excited quasiparticles

$$\rightarrow \frac{\partial S_s}{\partial T} > \frac{\partial S_n}{\partial T} \Rightarrow C_s \text{ is getting larger than } C_n$$



# 4.3 Thermodynamic Quantities

iii.  $T \simeq T_c$ :



$\Delta(T) \rightarrow 0 \rightarrow$  we can replace  $E_{\mathbf{k}}$  by  $|\xi_{\mathbf{k}}|$ :

$$C_s = \underbrace{\frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(\xi_{\mathbf{k}})}{\partial \xi_{\mathbf{k}}}}_{\text{normal state specific heat}} \left( \xi_{\mathbf{k}}^2 - \frac{1}{2} T \frac{d\Delta^2(T)}{dT} \right)$$

normal state specific heat  $C_n = \frac{\pi^2}{3} D(E_F) k_B^2 T$

finite for  $T < T_c$   
zero for  $T > T_c$

} *jump of specific heat*



M. A. Biondi *et al.*,  
Rev. Mod. Phys. **30**, 1109-1136 (1958)

# 4.3 Thermodynamic Quantities

iii.  $T \simeq T_c$ : *jump of specific heat* (we can replace  $E_{\mathbf{k}}$  by  $|\xi_{\mathbf{k}}|$ )

$$\Delta C = (C_s - C_n)_{T=T_c} = \frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(\xi_{\mathbf{k}})}{\partial \xi_{\mathbf{k}}} \left( -\frac{1}{2} T \frac{d\Delta^2(T)}{dT} \right)_{T=T_c}$$

$$\Delta C = D(E_F) \left( -\frac{d\Delta^2(T)}{dT} \right)_{T=T_c} \underbrace{\int_{-\infty}^{\infty} -\frac{\partial f(\xi_{\mathbf{k}})}{\partial \xi_{\mathbf{k}}} d\xi_{\mathbf{k}}}_{=1}$$

we use  $\Delta(T) \simeq 1.74 \left(1 - \frac{T}{T_c}\right)^{1/2}$  for  $T$  close to  $T_c$  and  $\Delta(0) = 1.76 k_B T_c$  and obtain

$$\Delta C \simeq 4.7 D(E_F) k_B^2 T_c$$

with  $C_n(T_c) = \frac{\pi^2}{3} D(E_F) k_B^2 T_c = \gamma T_c$  we finally obtain

$$\left( \frac{\Delta C}{C_n} \right)_{T=T_c} \simeq \frac{4.7}{\pi^2/3} = 1.43$$


**further key prediction of BCS theory**  
(in good agreement with experiment)

result from phenomenological treatment: **(Rutgers formula)**


$$\left( \frac{\Delta C}{C_n} \right)_{T=T_c} = \frac{1}{C_n} \frac{T_c}{\mu_0} \left( \frac{\partial B_{\text{cth}}}{\partial T} \right)_{T=T_c}^2 = \frac{1}{C_n} \frac{8}{T_c} \frac{B_{\text{cth}}^2(0)}{2\mu_0} = \frac{8}{\gamma T_c^2} \frac{B_{\text{cth}}^2(0)}{2\mu_0} = \frac{6}{\pi^2} \left( \frac{\Delta(0)}{k_B T_c} \right)^2 = \frac{6}{\pi^2} (1.76)^2 = 1.88$$

difference comes from  $B_{\text{cth}}(T)$

# 4.3 Thermodynamic Quantities



N.E. Phillips,  
Phys. Rev. **114**, 676 (1959)



M. A. Biondi et al.,  
Rev. Mod. Phys. **30**, 1109-1136 (1958)



# Summary of Lecture No. 9 (1)

- Bogoliubov-Valatin transformation → BCS Gap Equation and Excitation Spectrum**

$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \sum_{\mathbf{k}} \left\{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* + \underbrace{(\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow})}_{\mathcal{B}_{\mathbf{k}}^\dagger} \underbrace{\mathcal{U}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}^\dagger}_{\varepsilon_{\mathbf{k}}} \begin{pmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}} \\ -\Delta_{\mathbf{k}}^* & -\xi_{\mathbf{k}} \end{pmatrix} \underbrace{\mathcal{U}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}^\dagger}_{\mathcal{B}_{\mathbf{k}}} \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix} \right\} = \sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^\dagger + \mathcal{B}_{\mathbf{k}}^\dagger \mathcal{U}_{\mathbf{k}}^\dagger \varepsilon_{\mathbf{k}} \mathcal{U}_{\mathbf{k}} \mathcal{B}_{\mathbf{k}} \}$$

$$\mathcal{B}_{\mathbf{k}}^\dagger = (\alpha_{\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}) = (\hat{c}_{\mathbf{k}\uparrow}^\dagger, \hat{c}_{-\mathbf{k}\downarrow}) \mathcal{U}_{\mathbf{k}} \quad \mathcal{B}_{\mathbf{k}} = (\alpha_{\mathbf{k}}, \beta_{-\mathbf{k}}^\dagger) = \mathcal{U}_{\mathbf{k}}^\dagger \begin{pmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{pmatrix}$$

**Bogoliubov quasiparticles:** → superposition of electron and hole states with opposite momentum and spin

Task: find unitary matrix ( $\mathcal{U}_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}^\dagger = 1$ ) that makes the transformed energy matrix  $\tilde{\varepsilon}_{\mathbf{k}} = \mathcal{U}_{\mathbf{k}}^\dagger \varepsilon_{\mathbf{k}} \mathcal{U}_{\mathbf{k}}$  diagonal:

$$\mathcal{U}_{\mathbf{k}} = \begin{pmatrix} u_{\mathbf{k}} & v_{\mathbf{k}}^* \\ -v_{\mathbf{k}} & u_{\mathbf{k}}^* \end{pmatrix} \quad \mathcal{U}_{\mathbf{k}}^\dagger = \begin{pmatrix} u_{\mathbf{k}}^* & -v_{\mathbf{k}}^* \\ v_{\mathbf{k}} & u_{\mathbf{k}} \end{pmatrix} \quad \Rightarrow \quad \mathcal{U}_{\mathbf{k}}^\dagger \varepsilon_{\mathbf{k}} \mathcal{U}_{\mathbf{k}} = \begin{pmatrix} E_{\mathbf{k}} & D_{\mathbf{k}} \\ -D_{\mathbf{k}} & -E_{\mathbf{k}} \end{pmatrix} \quad \Rightarrow \quad D_{\mathbf{k}} = 0 \quad E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

- reformulation of the BCS Hamilton operator**

$$\mathcal{H}_{\text{BCS}} - \mathcal{N}\mu = \underbrace{\sum_{\mathbf{k}} \{ \xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta_{\mathbf{k}} g_{\mathbf{k}}^* \}}_{\text{mean-field contribution}} + \underbrace{\sum_{\mathbf{k}} \{ E_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + E_{\mathbf{k}} \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}} \}}_{\text{contribution of spinless Fermion system}}$$

→ minimization of free energy yields BCS gap equation for finite  $T$

**mean-field contribution**  
differs from the normal state value by the condensation energy (see below)

contribution of **spinless Fermion system** with two kind of **quasiparticles** described by operators  $\alpha_{\mathbf{k}}^\dagger, \alpha_{\mathbf{k}}$  and  $\beta_{-\mathbf{k}}^\dagger, \beta_{-\mathbf{k}}$  and excitation energies  $\pm E_{\mathbf{k}}$   
spinless **quasiparticles** since they consist of superposition of spin- $\uparrow$  and spin- $\downarrow$  electrons

# Summary of Lecture No. 9 (2)

- *minimization of free energy yields BCS gap equation:*

$$\Delta_{\mathbf{k}}^* = - \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}^*}{2E_{\mathbf{k}'}} \tanh \left( \frac{E_{\mathbf{k}'}}{2k_B T} \right) \quad \text{BCS gap equation}$$

- *analytical solution with simplifications:*

$$V_{\mathbf{k},\mathbf{k}'} = -V_0, \Delta_{\mathbf{k}} = \Delta, V_0 D(E_F) \ll 1: \text{weak coupling approximation} \rightarrow 1 = V_0 \sum_{\mathbf{k}'} \frac{1}{2E_{\mathbf{k}'}} \tanh \left( \frac{E_{\mathbf{k}'}}{2k_B T} \right)$$

$$T \ll T_c$$

$$\Delta(0) \simeq 2\hbar\omega_D e^{-\frac{2}{V_0 D(E_F)}}$$

$$T \simeq T_c$$

$$k_B T_c = 1.13 \hbar\omega_D e^{-\frac{2}{V_0 D(E_F)}}$$



$$\frac{\Delta(0)}{k_B T_c} = \frac{\pi}{e^\gamma} = 1.764$$

- *condensation energy at  $T = 0$*

$$V_{\mathbf{k},\mathbf{k}'} = -V_0, \Delta_{\mathbf{k}} = \Delta, V_0 D(E_F) \ll 1: \text{weak coupling approximation}$$

$$E_{\text{kond}}(0) = \langle \mathcal{H}_{\text{BCS}} \rangle - \langle \mathcal{H}_n \rangle = -D(E_F) \Delta^2(0) / 4$$


comparison to  $E_{\text{cond}}(0) = -B_{\text{cth}}^2(0) / 2\mu_0$  (thermodynamics) yields

$$B_{\text{cth}}(0) = \sqrt{\frac{\mu_0 D(E_F) \Delta^2(0)}{2V}}$$

# Summary of Lecture No. 9 (3)

- density of states:**

$$D_s(E_{\mathbf{k}}) = D_n(\xi_{\mathbf{k}}) \frac{d\xi_{\mathbf{k}}}{dE_{\mathbf{k}}} = \begin{cases} D_n(E_F) \frac{E_{\mathbf{k}}}{\sqrt{E_{\mathbf{k}}^2 - \Delta^2}} & \text{for } E_{\mathbf{k}} > \Delta \\ 0 & \text{for } E_{\mathbf{k}} < \Delta \end{cases}$$



$$E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

- BCS prediction for thermodynamic quantities**

$$S_s = -2k_B \sum_{\mathbf{k}} \left\{ \underbrace{[1 - f(E_{\mathbf{k}})] \ln[1 - f(E_{\mathbf{k}})]}_{\text{hole like}} + \underbrace{f(E_{\mathbf{k}}) \ln[f(E_{\mathbf{k}})]}_{\text{electron like}} \right\}$$

entropy


$$S = -k_B \sum_n p_n \ln p_n$$

$$C_s = \frac{2}{T} \sum_{\mathbf{k}} -\frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \left( E_{\mathbf{k}}^2 - \frac{1}{2} T \frac{d\Delta_{\mathbf{k}}^2(T)}{dT} \right)$$

heat capacity

⇒  $C_s \propto T^{-\frac{3}{2}} e^{-\frac{\Delta(0)}{k_B T}} \propto T^{-\frac{3}{2}} e^{-1.76 \frac{T_c}{T}} @ T \ll T_c$

exponential decrease of heat capacity at low T





Walther  
Meißner  
Institut



BAYERISCHE  
AKADEMIE  
DER  
WISSENSCHAFTEN

Technische  
Universität  
München



# Superconductivity and Low Temperature Physics I



**Lecture No. 10**

**R. Gross**

**© Walther-Meißner-Institut**

## 4. Microscopic Theory

### 4.1 Attractive Electron-Electron Interaction

#### 4.1.1 Phonon Mediated Interaction

#### 4.1.2 Cooper Pairs

#### 4.1.3 Symmetry of Pair Wavefunction

### 4.2 BCS Ground State

#### 4.2.1 The BCS Gap Equation

#### 4.2.2 Ground State Energy

#### 4.2.3 Bogoliubov-Valatin Transformation and Quasiparticle Excitations

### 4.3 Thermodynamic Quantities



### 4.4 Determination of the Energy Gap

#### 4.4.1 Specific Heat

#### 4.4.2 Tunneling Spectroscopy

### 4.5 Coherence Effects

# 4.4 Determination of the Energy Gap

- energy gap determines excitation spectrum of superconductors
  - we can use quantities that depend on excitation spectrum to determine  $\Delta$ 
    1. specific heat
    2. tunneling conductance
    3. microwave and infrared absorption
    4. ultrasound attenuation
    5. ....
- we concentrate on tunneling spectroscopy in the following (specific heat already discussed in previous subsection)

# 4.4.2 Tunneling Spectroscopy


tunneling of quasiparticle excitations between two superconductors separated by thin tunneling barrier

- SIS tunnel junction:




- fabrication by thin film technology and patterning techniques


by shadow masks ( $\approx$  mm)  
 by optical lithography ( $\approx$   $\mu$ m)  
 by e-beam lithography ( $\approx$  10 nm)



- sketch:



top view:



# 4.4.2 Tunneling Spectroscopy

- tunneling processes result in finite coupling of SC 1 and SC 2, described by tunneling hamiltonian

$$\mathcal{H}_{\text{tun}} = \sum_{\mathbf{k}\mathbf{q}\sigma} T_{\mathbf{k}\mathbf{q}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{q}\sigma} + c.c.$$

tunnel matrix element


describes the creation of electron  $|\mathbf{k}\sigma\rangle$  in one SC and the annihilation of electron  $|\mathbf{q}\sigma\rangle$  in the other

- tunneling into state  $|\mathbf{k}\sigma\rangle$  only possible if pair state  $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$  is empty

→ resulting tunneling probability is  $\propto |u_{\mathbf{k}}|^2 |T_{\mathbf{k}\mathbf{q}}|^2$

- for each state  $|\mathbf{k}\sigma\rangle$  there exists a state  $|\mathbf{k}'\sigma\rangle$  with  $E_{\mathbf{k}} = E_{\mathbf{k}'}$  but with  $\xi_{\mathbf{k}'} = -\xi_{\mathbf{k}}$

→ resulting tunneling probability is  $\propto |u_{\mathbf{k}'}|^2 |T_{\mathbf{k}'\mathbf{q}}|^2 \stackrel{=}{=} |v_{\mathbf{k}}|^2 |T_{\mathbf{k}\mathbf{q}}|^2$   
 $|u(-\xi_{\mathbf{k}})| = |v(\xi_{\mathbf{k}})|$



➡ total tunneling probability  $\propto (|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2) |T_{\mathbf{k}\mathbf{q}}|^2 = |T_{\mathbf{k}\mathbf{q}}|^2$  **does not depend on coherence factors**

→ *simple „semiconductor model“ for quasiparticle tunneling is applicable*



# 4.4.2 Tunneling Spectroscopy

elastic tunneling between two metals (NIN):

$$I_{1 \rightarrow 2} = C \int_{-\infty}^{\infty} |T|^2 \underbrace{D_1(E)f(E)}_{\text{occupied states in } N_1} \underbrace{D_2(E + eV) [1 - f(E + eV)]}_{\text{empty states in } N_2} dE$$


- net tunneling current:

$$I_{nn}(V) = C \int_{-\infty}^{\infty} |T|^2 D_1(E) D_2(E + eV) [f(E) - f(E + eV)] dE$$

- for  $eV \ll \mu$  and  $\mu \simeq E_F$  we can use  $D_n(E + eV) \simeq D_n(E_F) = \text{const.}$

$$I_{nn}(V) = C |T|^2 D_{n1}(E_F) D_{n2}(E_F) \underbrace{\int_{-\infty}^{\infty} [f(E) - f(E + eV)] dE}_{=eV}$$

$$I_{nn}(V) = C |T|^2 D_{n1}(E_F) D_{n2}(E_F) e V = G_{nn} V$$



# 4.4.2 Tunneling Spectroscopy

elastic tunneling between N and S (NIS junction):



$$I_{ns}(V) = \underbrace{C|T|^2 D_{n1}(E_F) D_{n2}(E_F)}_{=G_{nn}/e} \int_{-\infty}^{\infty} \frac{D_{s2}(E)}{D_{n2}(E_F)} [f(E) - f(E + eV)] dE$$

$$I_{ns}(V) = \frac{G_{nn}}{e} \int_{-\infty}^{\infty} \frac{D_{s2}(E)}{D_{n2}(E_F)} [f(E) - f(E + eV)] dE$$



- analytical solution for  $T = 0$

$$I_{ns}(V) = \frac{G_{nn}}{e} \int_{-\infty}^{\infty} \frac{|E|}{|E^2 - \Delta^2|^{1/2}} [f(E) - f(E + eV)] dE$$

$$I_{ns}(V) = \begin{cases} 0 & |eV| < \Delta \\ \frac{G_{nn}}{e} [(eV)^2 - \Delta^2]^{1/2} & |eV| \geq \Delta \end{cases}$$



# 4.4.2 Tunneling Spectroscopy




# 4.4.2 Tunneling Spectroscopy

## differential tunneling conductance of NIS junction

$$I_{ns}(V) = \frac{G_{nn}}{e} \int_{-\infty}^{\infty} \frac{D_{s2}(E)}{D_{n2}(E_F)} [f(E) - f(E + eV)] dE$$

$$G_{ns}(V) = \frac{\partial I_{ns}(V)}{\partial V} = G_{nn} \int_{-\infty}^{\infty} \frac{D_{s2}(E)}{D_{n2}(E_F)} \underbrace{\left[ -\frac{\partial f(E + eV)}{\partial (eV)} \right]}_{\text{Bell-shaped weighting function}} dE \quad dE = e dV$$

Bell-shaped weighting function with width  $\approx 4k_B T$  peaked at  $E = eV$   
 $\rightarrow$  approaches  $\delta$ -function for  $T \rightarrow 0$


  $G_{ns}(V) = G_{nn} \frac{D_{s2}(eV)}{D_{n2}(E_F)} \quad @ T = 0$       **measurement of  $G_{ns}(V)$  allows determination of  $D_{s2}(eV)$  and  $\Delta$ , for  $T > 0$ ,  $G_{ns}(V)$  measures DOS smeared out by  $\pm k_B T$**

- at  $T > 0$ : finite conductance at  $eV \ll \Delta$  due to smeared Fermi distribution, calculation yields

$$\left. \frac{G_{ns}}{G_{nn}} \right|_{eV \ll \Delta} = \left( \frac{2\pi\Delta}{k_B T} \right) e^{-\Delta/k_B T}$$

**exponential  $T$ -dependence can be used for temperature measurement, particle detectors, ...**

# 4.4.2 Tunneling Spectroscopy




I. Giaever, K. Megerle,  
 Phys. Rev. **122**, 1101-1111 (1961)


# 4.4.2 Tunneling Spectroscopy

elastic tunneling between two superconductors: SIS junction

$$I_{SS}(V) = \frac{G_{nn}}{e} \int_{-\infty}^{\infty} \frac{D_{S1}(E + eV)}{D_{n1}(E_F)} \frac{D_{S2}(E)}{D_{n2}(E_F)} [f(E) - f(E + eV)] dE$$




- strong increase of  $I_{SS}$  @  $eV = 2\Delta$
- $I_{SS} \rightarrow I_{nn}$  @  $eV \gg 2\Delta$
- $I_{SS} \propto e^{-\Delta/k_B T}$  @  $eV < 2\Delta$



# 4.4.2 Tunneling Spectroscopy

interpretation of tunneling in SIS junction at  $T = 0$

- single electron tunnels from left to right:




- energy balance:  $-E_F^{(\text{left})} + E_{\mathbf{k}}^{(\text{left})} + E_F^{(\text{right})} + E_{\mathbf{k}}^{(\text{right})}$   
 $e^-$  moves from left to right      generation of two qp

$$E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}$$

- required voltage:  $eV = E_{\mathbf{k}}^{(\text{left})} + E_{\mathbf{k}}^{(\text{right})}$
- minimal voltage:  $eV = \Delta_1 + \Delta_2 = 2\Delta$  for  $\Delta_1 = \Delta_2$

# 4.4.2 Tunneling Spectroscopy

current-voltage characteristics of SIS junction at finite temperatures






# 4.4.2 Tunneling Spectroscopy

special case: SIS tunnel junction with  $\Delta_1 \neq \Delta_2$

- at  $eV = \Delta_2 - \Delta_1$  the two singularities in the DOS are facing each other
  - $\Rightarrow$  maximum of the tunneling current
  - $\Rightarrow$  negative differential resistance



# 4.5 Coherence Effects

- description of an external perturbation on the electrons in a metal

$$\mathcal{H}_1 = \sum_{\mathbf{k}\sigma, \mathbf{k}'\sigma'} P_{\mathbf{k}'\sigma', \mathbf{k}\sigma} c_{\mathbf{k}'\sigma'}^\dagger c_{\mathbf{k}\sigma}$$

interaction hamiltonian

$|P_{\mathbf{k}'\sigma', \mathbf{k}\sigma}|^2$  corresponds to transition probability

- description of the external perturbation on the electrons in a superconductor

→ more complicated since there is a coherent superposition of occupied one-electron states

$$\left. \begin{aligned} \hat{c}_{\mathbf{k}\uparrow}^\dagger &= u_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + v_{\mathbf{k}} \beta_{-\mathbf{k}} \\ \hat{c}_{-\mathbf{k}\downarrow} &= -v_{\mathbf{k}}^* \alpha_{\mathbf{k}}^\dagger + u_{\mathbf{k}} \beta_{-\mathbf{k}} \\ \hat{c}_{\mathbf{k}\uparrow} &= u_{\mathbf{k}} \alpha_{\mathbf{k}} + v_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger &= -v_{\mathbf{k}} \alpha_{\mathbf{k}} + u_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger \end{aligned} \right\} \begin{aligned} c_{\mathbf{k}'\uparrow}^\dagger c_{\mathbf{k}\uparrow} &= (u_{\mathbf{k}'}^* \alpha_{\mathbf{k}'}^\dagger + v_{\mathbf{k}'} \beta_{-\mathbf{k}'}^\dagger) (u_{\mathbf{k}} \alpha_{\mathbf{k}} + v_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger) \\ c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} &= (-v_{\mathbf{k}} \alpha_{\mathbf{k}} + u_{\mathbf{k}}^* \beta_{-\mathbf{k}}^\dagger) (-v_{\mathbf{k}'}^* \alpha_{\mathbf{k}'}^\dagger + u_{\mathbf{k}'} \beta_{-\mathbf{k}'}^\dagger) \end{aligned} \quad \text{connect the same qp states}$$

→ matrix elements  $|P_{\mathbf{k}'\sigma', \mathbf{k}\sigma}|^2$  have to be multiplied by so-called coherence factors

$(u_{\mathbf{k}} u_{\mathbf{k}'} \mp v_{\mathbf{k}} v_{\mathbf{k}'})^2$  for scattering of quasiparticles

$(v_{\mathbf{k}} u_{\mathbf{k}'} \pm u_{\mathbf{k}} v_{\mathbf{k}'})^2$  for creation or annihilation of quasiparticles

$u_{\mathbf{k}}, v_{\mathbf{k}}$  are assumed real


see e.g.

M. Tinkham


Introduction to Superconductivity

# 4.5 Coherence Effects


temperature dependence of low-frequency absorption processes in superconductors



# 4.5 Coherence Effects



Ultrasound Attenuation in  $Sr_2RuO_4$ :  
 An Angle-Resolved Study of the Superconducting Gap Function  
 C. Lupien, W. A. MacFarlane, Cyril Proust, Louis Taillefer, Z. Q. Mao, and Y. Maeno  
 Phys. Rev. Lett. **86**, 5986 (2001)



A.G. Redfield, *Nuclear Spin Relaxation Time in Superconducting Aluminum*.  
 Phys. Rev. Lett. **3**, 85–86 (1959)  
 L.C. Hebel, *Theory of Nuclear Spin Relaxation in Superconductors*.  
 Phys. Rev. **116**, 79–81 (1959).